



Paper number 115

ENVIRONMENT DEPARTMENT PAPERS

Climate Change Series

Transitioning to Climate Resilient Development

Perspectives from Communities in Peru

Frank Sperling, with
Corinne Valdivia, Roberto Quiroz, Roberto Valdivia,
Lenkiza Angulo, Anton Seimon, and Ian Noble

May 2008



Sustainable Development Vice Presidency



THE WORLD BANK ENVIRONMENT DEPARTMENT

Transitioning to Climate Resilient Development

Perspectives from Communities in Peru

Frank Sperling

with

Corinne Valdivia

Roberto Quiroz

Roberto Valdivia

Lenkiza Angulo

Anton Seimon

Ian Noble

May 2008

© The International Bank for Reconstruction
and Development/THE WORLD BANK
1818 H Street, N.W.
Washington, D.C. 20433, U.S.A.

Manufactured in the United States of America
First printing May 2008

The findings, interpretation, and conclusions presented in this report are those of the author and should not be attributed to the World Bank, its affiliated organizations, or to members of its Board of Executive Directors or the countries they represent. The presentation of material in this document and the geographical designations employed do not imply expression of any opinion on the part of the World Bank concerning the legal status of a country, territory or area, or concerning the delimitation of its boundaries.

Design: Jim Cantrell
Cover photos: Frank Sperling

Contents

ACKNOWLEDGMENTS	vii
EXECUTIVE SUMMARY	1
INTRODUCTION — CLIMATE RESILIENT DEVELOPMENT	5
<i>Chapter 1: Climate Resilient Development — A Country Perspective</i>	9
Disaster Risk — An Overview	11
El Niño’s Influence on Peru’s Climate	11
<i>Forecasting El Niño</i>	14
Institutional Structures for Disaster and Climate Risk Management	16
<i>Climate Change — Challenge and Institutional Response</i>	18
<i>Climate Change and El Niño</i>	19
<i>Chapter 2: Climate Resilient Development — Community Perspectives</i>	21
Approach	21
Perspectives from Communities in Piura and Puno	25
Disaster Risks and Impacts	25
<i>Piura</i>	25
<i>Puno</i>	29
<i>Spatial Disaster Risk</i>	31
<i>Temporal Exposure to Climatic Risks</i>	32
<i>Impacts</i>	34
<i>Comparison of Risk Exposure of Communities in Piura and Puno</i>	36
Vulnerability	36
<i>Socioeconomic Vulnerability</i>	37
<i>Environmental Vulnerability</i>	38
<i>Long-term Consequences of Vulnerability</i>	38
<i>Key Aspects of Vulnerability</i>	38
Coping and Adaptation Strategies	39
<i>Coping Strategies</i>	40
<i>Adaptation Practices</i>	41
<i>External Assistance and Barriers</i>	43
External Information—Weather Forecasts, Early Warning, and Seasonal Climate Outlooks	44
<i>Access to and Use of Information in Piura</i>	45
<i>Weather and Climate Information in Puno</i>	47
<i>Reflections on Community Responses</i>	48
Use of Local Knowledge in the Prediction of Climatic and Environmental Changes	50

<i>Piura</i>	50
<i>Puno</i>	50
<i>Gauging the Role and Validity of Indicators</i>	56
Perceptions of Changing Climate Conditions	57
<i>Piura: Perceived Climatic Changes in Northwestern Peru</i>	57
<i>Puno: Perceived Climatic Changes in the Peruvian Altiplano</i>	57
<i>Are These Perceptions True?</i>	58
Piura	58
Puno	60

Chapter 3: Conclusions and Recommendations 65

Use of Climate Information	66
<i>Research</i>	66
Environmental Risks and Change	67
Climate Forecasts and Local Indicators	67
Agriculture and Livestock Management and Market Research	67
Incentive Systems	67
Tracking Progress	68
<i>Community Action on Climate Risk Management</i>	68
Awareness Building	68
Community-Level Investments	69
<i>Enabling Frameworks: Institutional Coordination and Support</i>	70

REFERENCES 73

APPENDIXES

<i>Appendix A: Spatial Hazard Maps of Puno Communities</i>	77
<i>Appendix B: Warm (El Niño) and Cold (La Niña) Phases of ENSO</i>	83
<i>Appendix C: Precipitation Totals in Relation to Seasonal SST Anomalies in Esperanza</i>	85
<i>Appendix D: Community-Specific Recommendations</i>	89
Parachique	89
<i>Measures that Do Not Require Outside Assistance</i>	89
<i>Measures that Require Outside Assistance</i>	89
Chato Grande	90
<i>Measures the Community Could Take without Outside Assistance</i>	90
<i>Measures that Require Outside Assistance</i>	90
Locuto	91
<i>Measures the Community Could Take without External Assistance (according to July 13 workshop)</i>	91
<i>Measures that Require Outside Assistance (according to July 13 workshop)</i>	92
<i>Measures that Could Be Implemented by the Community without External Assistance (part of the development plan for the left bank of the Piura River, in Tambogrande)</i>	92
<i>Measures that Require External Assistance (part of the development plan for the left bank of the Piura River, in Tambogrande)</i>	93
Cabuyal	94
<i>Measures the Community Could Take without Outside Assistance</i>	94
<i>Measures that Require Outside Assistance</i>	94

Appendix E: Project Proposal for Follow-Up Activities in the Altiplano 97

Background	97
Development Objective	97
Expected Contribution to Development Outcomes/Results	98
Expected Outputs and Results	98
Key Performance Indicators	98

Appendix F: Managing Climate Risks in Coastal Ecuador — A Brief Comparison of Community Perspectives 101**FIGURES**

1. Monthly Precipitation for (A) La Esperanza, and (B) Huancabamba	12
2. Piura, Northern Peru	13
3. Monthly Precipitation Averages for Puno from 1931–73	14
4. The NINO Regions	14
5. NINO 3.4 Forecasts: Lead Time by Month and Predictive Skill	15
6. Diagram of Information Flow through ERFEN Network in Peru	17
7. NINO 3 Region Climate Trends	20
8. Spatial Exposure to Hydrometeorological Hazards of the Community of Santa Maria	31
9. Time Line of Major Natural Hazard Events Recalled by the Communities	32
10. A Comparison of the Seasonal Precipitation Total from January to March (JFM) for La Esperanza with JFM SST Anomalies in the NINO 3.4 Region	48
11. Comparison between January Precipitation in La Esperanza and SST Values of the NINO 1.2 Region	49
12. Average (A) Annual, (B) Monthly Maximum, and (C) Monthly Minimum Temperatures for Esperanza, Northwestern Coastal Peru	59
13. Average Monthly (A) Maximum and (B) Minimum Temperatures for Huancabamba, Sierras, Upper Piura Watershed	61
14. Long-term Temperature Record for (A) Minimum, (B) Maximum, and (C) Average Temperature in Cusco, Peru	62
15. Precipitation Averaged over 30-year Periods for January to April for the Peruvian Altiplano Region	63

MAPS

1. The Country of Peru	10
2. Location of Communities Assessed in Piura	22
3. Location of Communities in Puno	23

TABLES

1. Annual GDP Growth Rates in Percent by Year, Peru	10
---	----

2.	Disasters in Peru (Recorded by District from 2000–04)	11
3.	Three-Month Temperature Anomalies for Classification of El Niño (EN) and La Niña (LN) Conditions, in Degrees Celsius	15
4.	Members of ENFEN–Peru	17
5.	Community Characteristics, Piura	22
6.	Community Characteristics, Puno	24
7.	Disaster Events, as Recalled and Ranked by Communities in Piura, Peru	26
8.	Impacts of Climate Hazards Identified in the Community Assessments in Piura	27
9.	Qualitative Ranking of Natural Hazards by Communities in Puno	30
10.	Livelihood Impacts of Drought and Floods, Communities of Puno	35
11.	Livelihood Impacts of Hail and Frost, Communities of Puno	36
12.	Conditions of Vulnerability Highlighted by Communities	39
13.	Destinations of Push Migration, Piura Communities	41
14.	Access to Weather and Climate Information	46
15.	Awareness, Use, and Trust of El Niño Information in Six Puno Communities, 2005 Workshop	47
16.	Indicators Used by Communities to Predict Weather Extremes and Climatic Conditions	51

Acknowledgments

This report is an output of the project Adaptation Strategies to the Environmental and Socioeconomic Impacts of El Niño for Rural Communities in Ecuador and Peru supported by the Trust Fund for Environmental and Socially Sustainable Development (TFESSD) and led by the Climate Change Team of the World Bank's Environment Department.

This report has been authored by Frank Sperling. It draws upon two major consultancy studies in Piura, northwestern Peru and in Puno, the region of the Peruvian altiplano, which were carried out by the Intermediate Technology Development Group (ITDG) and Centro de Investigacion de Recursos Naturales y Medio Ambiente (Research Center for Natural Resources and Environment; CIRNMA) under the supervision and guidance of the Centro Internacional de la Papa (International Potato Center; CIP) in collaboration with the University of Missouri. The author would especially like to acknowledge the contributions provided by Roberto Quiroz (CIP), Corinne Valdivia (University of Missouri), Roberto Valdivia (CIRNMA), and Lenkiza Angulo (ITDG). Their knowledge and insights were of great importance in describing the situation of the communities in coping and adapting to climatic risks. In addition, Anton Seimon of Columbia University provided advice and shared valuable information, which is gratefully recognized. The author would also like to thank Ian Noble, the Task Team Leader of the Project, for this support and ideas.

The report would not have been possible without the participation of the communities of Parachique, Chato Grande, Locuto, and Cabuyal in the Department of Piura and the communities of Santa Maria, Ancacca, Yanamocco, Candile, Accopata, and Alto Achuyo in the Department of Puno. Their willingness and openness for sharing their perspectives on vulnerability to climate risk, coping and adaptive capacities, institutional support, and existing barriers are gratefully acknowledged.

Furthermore, our gratitude is extended to the institutions working on climate and disaster risk management issues in Peru, who shared information and provide insights about the challenges Peru is facing through the climatic changes associated with El Niño Southern Oscillation (ENSO) and global warming. Thanks are particularly due to Julio Garcia and staff at Consejo Nacional del Ambiente (National Council for the Environment; CONAM) for sharing data for this report. Also staff at Servicio Nacional de Meteorología e Hidrología (National Meteorology and Hydrology Service; SENAMHI), Instituto Nacional de Defensa Civil (National Institute of Civil Defense; INDECI), Instituto del Mar del Perú (Peruvian Marine Research Institute; IMARPE), and Ministerio de Agricultura (Ministry of Agriculture; MINAG) provided the team with helpful background information. Tony Barnston and colleagues at the International Research Institute of Climate and Society (IRI) at Columbia kindly shared information about their research.

The project team would also like to express its gratitude to Liliana Sánchez (Universidad Nacional Agraria la Molina) for organizing the workshops in Puno and assisting in the coordination of the assessments. Hugo Li Pun, Victor Mares, Oscar Hildago, Pili Byrne, and Martha Huanes (all CIP) organized and facilitated the workshops in Piura and Lima. We are most grateful for their help and time. We also like to thank Jorge de la Cruz for preparing the maps of the study region included in this report. Finally, Frank Sperling would like to thank Graciete Moura for her support.

This report is accompanied by a consultancy commissioned on early warning systems and institutional disaster prevention structures in Peru and Ecuador, which was jointly carried out by Senter for utvikling og miljø (Center for Development and the Environment, SUM), Center for International Climate

and Environmental Research (CICERO), and Norway, which led to the report *Seasonal Climate Forecasts and Community Responses: Summary of Experiences in Peru and Ecuador* by Jemima García-Godos, Desmond McNeill, Karen O'Brien, and Lars Otto Næss. The report is available upon request. Finally, Ajay Mathur, Walter Vergara, Laura Tlaiye and Renan Poveda are gratefully recognized as current (or former) World Bank staff who provided helpful general support and insights throughout different stages of this project initiative.

Further inquiries about this report or the project should be directed to the Climate Change Team (Environment Department, ENV) of the World Bank or directly to Ian Noble, inoble@worldbank.org or Frank Sperling, fsperling@worldbank.org.

Executive Summary

In the face of global environmental change, climate risk management is increasingly recognized for its essential role of ensuring the sustainability of development processes. Climate change is altering the exposure of developing countries to weather-related hazards, often exacerbating already existing vulnerabilities. Over recent decades, the worldwide occurrence of weather-related disasters and socioeconomic impacts have increased. This is linked to a variety of factors, including demographic, economic, and environmental trends. High economic losses associated with disasters and the importance of climate-sensitive sectors for the economic progress of many developing countries have strengthened the call for climate-proofing development processes by proactively managing current and future climate risks.

Peru is among the countries highly vulnerable to climatic variability and extremes. The Andean mountain range acts as formidable barrier to air masses and helped shape contrasting climatic zones. In addition, the El Niño Southern Oscillation (ENSO) exerts a strong influence on the climatic variability. The warm (El Niño) and cold (La Niña) phases of ENSO have been linked to the occurrence of climatic extremes in coastal Northwestern Peru and high plateau (altiplano) extending southward from Peru.

Climatic changes triggered by El Niño events have caused substantial economic losses. El Niño has been associated with above normal rainfall conditions and floods in the northwestern Peru and more frequent drought conditions in the altiplano. The strong 1997–98 El Niño event caused losses of 3.6 billion USD in

Peru. Given high levels of rural poverty, many impacts, however, reside outside mainstream economy affecting nonmonetary assets of rural people.

Understanding how to address the semiperiodic occurrence of climatic patterns associated with ENSO, as well as the multi-hazard environment of Peru, provide important insights into capacity-building needs to prepare for long-term climate change, while also being of immediate benefit. Within this context, the community assessments of this study compare information and capacity-building needs in Northern Peru, where El Niño exerts a dominating influence on climatic conditions, with the situation of communities in the Peruvian altiplano, where the effects of El Niño are superimposed on a generally high degree of climate variability.

The major challenge for communities in Northern Peru is to forge livelihoods out of a semiarid environment. El Niño events represent both a challenge and an opportunity when the additional rainfall amount turns from a relief to destruction in the short term, but to green forests in the medium term. Climate risk management in this region means adapting to and coping with either too much or too little water, but also taking advantage of the positive effects of climate variability.

The challenge for communities in the altiplano of Peru is to cope with a highly variable climate on an annual and interannual basis. While El Niño is often associated with drought conditions in the altiplano, the communities may be exposed to a range of climatic

extremes in any year. During the planting season, droughts, floods, hail, cold spells, and snow may impact livelihoods. This situation has made it very difficult for households to accumulate and sustain assets, reinforcing high levels of poverty.

While climate forecasts have the potential to support communities in improving their agricultural outputs, current communication formats are limited in their value, as they are not well integrated into the general knowledge networks and decision-making processes of the communities. Climate forecasts tend to be compared with biological and physical indicators of local environmental conditions. Forecasts are disseminated predominantly through the radio. Disagreement between the communities' interpretation of traditional indicators and the message of the forecast creates confusion, given that the one-way communication process of scientific forecasts does not provide opportunity for consultation or feedback. While forecasts are probabilistic, if this information is falsely interpreted it generates a general distrust toward using external information. The acceptance of seasonal climate forecasts and early warnings is furthermore dependent on who communicates the information, as government institutions are often not trusted, while radio is a trusted source.

Communities tend to recall El Niño events according to the magnitude of negative impacts they have caused. In coastal Ecuador and Northern Peru, the floods associated with strong 1982–83 and 1997–98 El Niño events are remembered. Weaker events are generally not linked by communities to disasters. This may in part be explained by the fact that additional moisture supplied during El Niño years to the semi-arid environment is, with the exception of the flooding events, largely beneficial for the communities, as it recharges aquifers and allows for the expansion of rain-fed agriculture and livestock. Therefore, in the perception of the communities of coastal Ecuador and Northern Peru, the return period of El Niño events differs from the scientific definition, as focus is

predominantly placed on the major events. This needs to be taken into account when issuing El Niño forecasts to communities.

El Niño events do not represent a core concern to communities in Southern Peru, as their effect is masked by the multi-hazard environment.

Communities relate El Niño events largely to impacts in Northern Peru and the coast and not to their own situation. While El Niño events have been shown to suppress moisture transport to the altiplano, and hence are associated with drought conditions, the continuous exposure to climatic extremes makes it difficult to recognize the specific impact of El Niño. It is therefore questionable whether an emphasis on strengthening awareness about El Niño would be useful; rather, climate information should reflect the interest of the communities in learning more about their disaster risks and the general strengthening of early-warning capabilities for floods and droughts as well as other hazards, and association with sea surface temperature (SST) forecasts in the Pacific.

Specific measures aimed at explaining, forecasting, and responding to climatic risk can further reduce climate-related development processes. A better understanding of spatial vulnerabilities to specific climatic hazards is needed. Climate forecasts and early-warning systems need to be linked to traditional knowledge systems and decision-making bodies.

The ability to cope with climatic hazards is dependent on assets and the ability to diversify income sources.

Particularly in the altiplano, the possession of livestock mitigates the impact of disasters. As access to credit is generally lacking and monetary assets are limited, the sale of livestock often provides the only opportunity to acquire medicines, seeds, and building materials after a disaster. Households that can resort to alternative sources of income, such as off-farm employment, are able to store forage for and protect their livestock, and can plant different crop varieties and adjust other agricultural practices to better cope

with climate extremes. Communal support systems and sharecropping can also play an important role in reducing the impact of climate-related hazards. In poorer households, where land quality is low and access to resources or communal support is limited, family members may be forced to migrate to urban areas or other regions in search for alternative income sources. Linkages built outside the community are essential because hazards are shared by a high proportion of families.

Development can improve the general resilience to climatic shocks, if the hazard risks are appropriately considered when addressing basic development needs. While communities are exposed to a range of climate-related hazards, vulnerabilities are often directly linked to basic development issues, such as the absence of health care facilities, inappropriate housing and infrastructure, limited access to markets, lack of educational and technical training opportunities, and weak institutional support. In this context, it is crucial that development activities, such as improving infrastructure, consider the local climatic conditions. In the altiplano, this means recognizing the exposure of development activities to extremely cold and hot,

as well as dry and wet, conditions. In Northern Peru, this means considering the effect of sporadically extremely wet conditions on infrastructure, human health, and agriculture during El Niño years, while strengthening the capacity of communities to maintain assets during the otherwise prevalent dry conditions. Training in crop and pest management for specific climatic conditions, strengthening access to credit for households or communities, and improving awareness about institutional support systems for emergency relief are also likely to significantly reduce climate-related vulnerabilities of community-level development processes.

Research activities need to be strengthened to support the design of climate-resilient development processes by responding to community-level training and capacity-building needs. Priority concerns are the validation of biological and physical indicators, evaluating the impact of environmental change, identifying effective entry points for scientific climate forecasts, evaluation of crop and livestock management options, exploring viable credit and insurance schemes, and the role of markets in fostering climate-resilient production and natural resources management.

Introduction — Climate Resilient Development

The international quest for eradicating poverty and sustainable development not only has to address tremendous demographic and socioeconomic challenges, but also needs to adapt to a changing global environment. Global warming will alter average climatic conditions in which development processes take place and affect the exposure to extreme events. The changes to the climate system are increasingly visible through their impacts on physical and biological systems around the globe (e.g. IPCC 2001a, b, IPCC 2007 a,b; Parmesan 2003, Stott et al. 2004, Emanuel 2005, and others). These changes occur against a backdrop of already existing vulnerabilities to natural disasters, which are particularly high in terms of loss of life and relative economic damages in developing countries (World Bank 2005). Climate change also interacts with other environmental changes, such as land degradation, habitat fragmentation, biodiversity loss, and air pollution (e.g., IPCC 2001, 2007). It is particularly the poor countries, where the livelihoods and economic growth depend on climate-sensitive sectors and adaptive capacity is low in terms of institutional and socioeconomic capital, that are at the receiving end of these changes.

In recognition of these problems, 10 bi- and multilateral agencies came together to jointly author the report *Poverty and Climate Change*, which emphasized the importance of integrating climate risk management into the development processes if sustainable progress toward and especially beyond the Millennium Development Goals (MDGs) is to be made (AfDB et al. 2003). The importance of an integrative approach of addressing climate change was recently reiterated

in the World Bank's Investment Framework for Clean Energy and Development (World Bank 2006), which also addresses adaptation to climate change. A first estimate suggests that around 40 percent of the development project portfolio is at significant risk due to climate change. As the fingerprints of climate change are increasingly becoming visible, there is clearly the need to understand what practical measures can be undertaken to improve the resilience to current and future climate risks.

The challenge lies in addressing the pressing development needs of today, without locking into rigid development paths that only promise short-term relief and not a sustainable solution to these problems, because of their inability to adapt to a dynamic environment.

The aim of adaptation to climate change is to reduce the exposure to climate hazards and/or the underlying vulnerabilities to climatic changes. In contrast to the necessary complementary efforts to mitigate the rate and magnitude of climate change through the reduction and sequestration of greenhouse gases, adaptation is less tangible. It requires different measures that depend on the scale, geography, and time horizon of the development sector targeted. Human health, agriculture and fisheries, forestry, and infrastructure are among the development sectors most vulnerable to climate change and associated impacts (IPCC 2007b), illustrating that practical efforts to adapt to climate change will need to cover a broad array of professional disciplines and cultures. This is further complicated by the variety of physical changes in the environment that are triggered

by climate change, including sea level rise, climatic extremes, changes in temperature, precipitation, and other effects. These changes are not uniformly distributed across the globe, but are associated with impacts that play out over different temporal and spatial scales.

The term climate-resilient development implies that adaptation to climate change should not be seen as a separate process, but a continuous and integrated one that addresses present and future climate risks. The starting point in this process is to identify and address existing vulnerabilities to current climate variability and change. Addressing existing adaptation “deficits” can provide important learning lessons for tackling climate change (Burton and van Aalst 2004), while also providing immediate benefits.

Some caveats apply. When addressing vulnerabilities to current climatic conditions, adaptation measures which are part of longer-term processes should be flexible so they are capable of being adjusted to further change. The knowledge of change already has implications as it means that our experiences may only be of limited use in addressing future climatic conditions (Sperling and Szekely 2005). Hence, addressing current vulnerabilities is the first step; only the beginning of a process that needs to be adjusted and reevaluated over time. But making this first step is crucial as it reflects a change in attitudes; a shift from coping (ex post) to prevention and vulnerability reduction (ex ante).

Scope and Rationale of Study

The paper focuses on the abilities of communities in Peru to cope with climate variability, drawing on the results of project financed by the Trust Fund for Environmental and Socially Sustainable Development (TFESSD). The climatic changes induced by the semiperiodic occurrence of El Niño and La Niña represent a learning opportunity of processes that influence adaptive capacities and could hence provide

important lessons for preparing for the challenges of climate change.

The northwestern coastal areas of Peru, as well as the altiplano in the South, are under the influence of the El Niño Southern Oscillation (ENSO). ENSO represents a natural ocean atmosphere phenomenon, which is associated with changes in SST along the equatorial Pacific and off the coast of Ecuador and Peru and fluctuations in the intertropical surface pressure pattern in the Pacific (and Indian) Ocean. The two extreme phases of ENSO, El Niño and La Niña, affect the climate of the tropical Pacific and beyond. Besides influencing average climatic conditions and climate variability, they have also been associated with the occurrence of extreme events, which have triggered disasters with macroeconomic impacts in Peru and elsewhere.

In recent years, the scientific capacity to forecast ENSO extremes and warn of impending El Niño events has increased considerably. This is complemented by a growing institutional emphasis on proactive disaster risk management, which aims to not only respond to but also prevent the occurrence of disasters through reduction of underlying vulnerability factors.

The semiperiodic occurrence of El Niño and La Niña events represents the opportunity to explore how these scientific and institutional advances have benefited vulnerable communities, such as the ones in Peru, in increasing their resilience to associated climatic extremes.

Specifically, the TFESSD study set out to capture the perspectives and opinions of communities concerning their vulnerability to climatic hazards, and the role of weather and climate information in relation to other nonformal information sources for anticipating and adapting to climatic extremes. The communities were also asked about perceived changes in their environment. Building on these discussions the communities then identified, first individually and then

jointly with institutional representatives, measures that need to be taken to strengthen their adaptive capacity to ENSO events and general climate variability.

This study uses a bottom-up approach. The livelihoods of communities in rural areas of Peru are predominantly dependent on climate-sensitive sectors such as agriculture. Given their close connection with the environment, the understanding of how these communities perceive risks and how they adapt to and cope with climate hazards is important for identifying entry points for efforts aimed at building resilience. In this context, it is assumed that the successful use of climate information will depend on appropriate information formats that fit the decision-making structure of communities as stakeholders and also have their trust. In connection with collecting information of community knowledge on environmental predictors of

weather and climatic conditions and recommendations for capacity-building needs, it is hoped that the report will provide valuable, initial guidance on which elements play an important role in strengthening the adaptive capacity of communities to climate variability and change.

The objective of Part I is to place climate variability and change into the broader development of Peru and outline risk management structures. This overview will then be contrasted with the perceptions of risks and vulnerabilities and coping and adaptation strategies at the community level described in Part II for Northwestern Peru and the altiplano, which are regions considered highly vulnerable to climate-related hazards. The report concludes by developing a set of overarching and locale-specific recommendations.

1 Climate Resilient Development— A Country Perspective

Peru is a country characterized by a diverse range of climates and an abundance of natural resources and environments. The Andes divide the country into three regions.

The semiarid to arid coastal area includes the capital of Lima and represents the region with the most industrialized and westernized urban centers. Agricultural activities are either under large-scale irrigation or, as is often the case of the more poor and vulnerable communities, dependent on seasonal river flows and highly variable rainfall.

About 50 percent of Peru's population of roughly 28 million lives in sierra, the highland plateau between the two ranges of the Andean mountains. Most Peruvians of indigenous origin live in the Sierras. Rural livelihoods still rely heavily on subsistence farming and traditional livestock rearing, such as llama and alpaca.

The eastern region of Peru is comprised of the Amazon lowlands, characterized by a tropical climate and rainforests. It is the region that is most sparsely populated.

Over the recent years, Peru has experienced a stable macroeconomic situation, which has been attributed to sound economic policies and a favorable external environment (World Bank 2006). Since 2002, GDP growth rates have been around 4 percent and higher. High levels of poverty still persist, however.

One in two Peruvians is poor (World Bank 2006). The rural populations and especially the indigenous people

are the most vulnerable with the least economic means. On the national level, 64 percent of indigenous people live below the poverty line, and almost 80 percent of the rural indigenous population is poor (World Bank 2006; see also Hall and Patrinos 2005). In contrast, 40 and 66 percent of the nonindigenous are poor at the national level and in rural areas, respectively.

While economic shocks more greatly affect the urban population, it is particularly the rural population that is most affected by natural disasters (World Bank 2006). A recent analysis of the world's disaster hotspots conducted by the World Bank and Columbia University ranked Peru among the 50 countries most vulnerable to natural disasters (World Bank 2005).

As will be discussed in the subsequent sections, climate-related hazards feature prominently in Peru's disaster risk profile, especially climatic extremes associated with the semiperiodic occurrence of El Niño events. The economic and structural losses associated with the 1982–83 El Niño were estimated to be 3.38 billion USD (6.2 percent of 1983 gross domestic product [GDP]), while the estimated loss of the 1997–98 El Niño amounted to approximately 3.5 billion USD, which represents about 3 percent of the 1998 GDP. 1998 represents a year characterized by a decline in GDP (Table 1). Given the existing vulnerability to interannual climate variability and the high dependency of livelihoods and economic activities on natural resources, the implications of climate change for the exposure to extreme events and the provision of

natural ecosystem services are becoming a growing concern. This is reflected by the recent recognition in

the Country Assistance Strategy of climate change as a threat to development processes.

Table 1. Annual GDP Growth Rates in Percent by Year, Peru

Year	1998	1999	2000	2001	2002	2003	2004	2005
GDP growth (%)	-0.7	0.9	3.0	0.2	5.2	3.9	5.2	6.4

Source: World Bank 2006.

Map 1. The Country of Peru



Disaster Risk — An Overview

Peru is exposed to a broad range of natural hazards. While emphasis is placed here on climate risks, it is important to recognize that the country is located in a tectonically active zone and hence large areas are also prone to geological hazards. Peru ranks highest among Latin American countries in numbers of lives lost and affected by natural disasters (Charveriat 2000).

There is evidence of increasing frequency of natural disasters over the short as well as long-term (World Bank 2006). Environmental degradation, urbanization, and demographic trends are contributing to increasing disaster risk (World Bank 2006), while global warming may increasingly affect the exposure to weather-related hazards in the region (e.g., IPCC 2007 a, b).

Peru is divided into 25 departments. This study focuses particularly on the departments of Piura and Puno. Both departments are characterized by high vulnerability to climatic risks.

Piura is situated in northwestern Peru and encapsulates the region most noted for its vulnerability to El Niño events. The department of Puno is part of the Andean altiplano, which stretches from 15° to 21° S, and is shared by Peru, Bolivia, and Chile. While ENSO also exerts its influence, the climate of the altiplano is generally characterized by high annual and interannual variability. Livelihoods in the region have to cope with a broad variety of extremes, including floods, droughts, cold spells, and hail events.

According to the National Institute for Civil Defense (INDECI), the Department of Puno in the altiplano ranks highest in five individual disaster categories, as

Table 2. Disasters in Peru (Recorded by District from 2000–04)

Type of Natural Disaster	Total	Piura	Puno	Max	Average
Volcanic Activity	1			1	1.0
Aluvium	11			3	1.8
Avalanches	6			2	1.5
Landslides	496	17	20	69	19.8
Hail	120	0	39	39	4.8
Frost	556	9	124	124	22.2
Huaycos	244	2	1	29	9.8
Floods	1328	30	185	185	53.1
Heavy rains	1241	4	147	147	49.6
Ocean swells	14	0	0	4	0.6
Snow	367	0	72	72	14.7
Drought	221	21	0	91	8.8
Earthquake	229	0	5	118	9.2
Thunderstorms	45	1	6	13	1.8
Strong winds	1806	125	108	386	72.2
Others	213	1	0	93	8.5
Total	6898	210	707	707	275.9

Source: INDECI 2005.

well as in total number of disasters among all Peruvian departments over the recent five-year period from 2000 to 2004 (Table 2), which represents a time frame characterized by the absence of a strong El Niño event.

El Niño's Influence on Peru's Climate

Originally, El Niño was perceived as a local phenomenon. Anecdotal evidence suggests that fishermen had observed warming of the usually cold coastal waters off the coast of Peru, which occurred semiperiodically around Christmastime and hence led to phenomenon being named after the child of Christ (e.g., Glantz 1996). Associated with these SST changes were the disappearance of fish, starvation of sea birds, and flooding events in Northern Peru and coastal Ecuador.

Building on the research of Sir Gilbert Walker and, later, Jacob Bjerkness, scientists began to reveal that the local changes associated with El Niño in Peru are linked to changes, which play out over much larger spatial scales.

El Niño is now recognized as one extreme outcome of atmospheric interactions between the ocean and atmosphere in the tropical Pacific, known today as the

El Niño Southern Oscillation (ENSO). The ENSO counterpart of El Niño is La Niña. Both phenomena are exerting a strong influence on the global climate, affecting regional temperature and precipitation patterns as well as extreme events.

During non-El Niño years, the atmospheric pressure in the eastern Southern Pacific (east of Tahiti) is higher than in the west (near Darwin, Australia). As El Niño conditions evolve, this pressure gradient weakens, turning off the engine fueling trade winds. The Intertropical Convergence Zone (ITCZ) is displaced southward while SSTs warm along the eastern equatorial Pacific and off the Peruvian coast. This results in rainfall anomalies in coastal regions of Northern Peru. Furthermore, the warming of the coastal waters and weakening of the winds chokes off the upwelling of cooler nutrient-rich waters, which leads to the migration of fish stocks.


Due to the ENSO phenomenon, Peru is exposed to large climatic interannual variations. The climatic changes associated with El Niño and La Niña, however, differ considerably between the coastal region of northwestern Peru and the altiplano in the south.

In northwestern Peru, which includes the

department of Piura, El Niño events are associated with a prolonged and intensified rainy season (Figure 1). While there is commonality in the direction of change, each El Niño event has its individual temporal and spatial characteristics.

While El Niño events are often accompanied by climatic extremes in northwestern Peru, it also has some

Figure 1. Monthly Precipitation for (A) La Esperanza, and (B) Huancabamba



Note: Monthly precipitation averages for (a) La Esperanza (latitude: 04°55'04", longitude: 81°03'38"; altitude: 12 meters [m] above sea level [a.s.l.]), 1960–2000, and (b) Huancabamba (05°15'; 79°43'; 1,952 m) 1970–96, Peru. La Esperanza, representing the coastal region of northwestern Peru, shows a clear increase in rainfall during El Niño years, in particular during the strong events of 1982–83 and 1997–98. El Niño years are also associated with spikes in precipitation in the sierras, as reflected in the rainfall record of Huancabamba, but rainfall in the highlands (alto) is generally more variable than in the low (bajo) and middle (medio) Piura.

Source: Authors' analysis, based on data provided by CONAM.

beneficial effects. The usually prevalent arid conditions limit agricultural activities to areas with (largely seasonal) water supply from the rivers or access to irrigation. The extra water input during El Niño years also increases visibly the ecosystem productivity of the region (Figure 2). El Niño should therefore not solely be associated with disasters, but its positive effects also have to be recognized.

In contrast to northwestern Peru, the altiplano, which includes the department of Puno, is exposed to high annual and interannual climate variability. A variety of extreme events, including droughts, cold spells, floods, hail, and unseasonable snowfall are frequently

experienced and represent a challenge to livelihoods based on agricultural practices (see Table 1).

The climate is generally semiarid and the rainy season largely confined to austral summer months. Substantial precipitation events usually begin around mid-December and last until April (Figure 3). As a result, agricultural activities are mainly limited to one planting season. While in Northwestern Peru rainfall totals are substantially lower, two to three annual planting cycles can be possible during years with abundant water supply due to the widespread use of irrigation.


Another general characteristic of the climate in the altiplano is a pronounced diurnal temperature range. The difference between daily maximum and minimum temperatures can be 35° C or more (Jordan 1991).

The effects of El Niño on the South American altiplano are more variable. El Niño conditions are associated with below-normal rainfall conditions or drought conditions, while La Niña appears to favor wetter conditions. The ENSO signals not as consistent as in northwestern Peru, however, and important exceptions occur.

El Niño conditions tend to suppress the moisture transport from Amazonian lowlands into the altiplano by limiting the number of days with easterly flow due to the associated warming of the lower atmosphere (Garreaud and Aceituno 2001). During La Niña years the opposite occurs. Increases in days with easterly winds promote the upslope transport of moist air into the altiplano, hence favoring wet conditions.


While consequently El Niño is associated with drought conditions and La Niña with floods, this is not always the case, as the ENSO influence is highly sensitive to positive and negative anomalies in the flow of easterly winds (Garreaud and Aceituno 2001). Hence, other local-to-mesoscale disturbances can support or offset the influence of ENSO on the climate of the altiplano. For example, the strong El Niño of 1982–83 led to

Figure 2. Piura, Northern Peru



Note: The satellite image shows (a) the revitalized vegetation in the coastal regions during the April 1998, an El Niño year, in comparison to (b) the prevalent dry environment during the same month in 2001, a non-El Niño year. Source: CIP.

Figure 3. Monthly Precipitation Averages for Puno from 1931–73



Note: Long-term averages of monthly precipitation totals with the associated standard deviations are presented for Puno. The monthly values were averaged over the time-frame from 1931–73, which represents a period of continuous data recording. Each month is based on between 40 to 43 values.

Source: Seimon.

widespread drought in the altiplano, while this was not the case during the strong El Niño of 1997–98.

Seimon (2004) suggests that ENSO may also influence the length of the seasons in the altiplano, with El Niño shortening and La Niña prolonging the summer (rainy) season. In general, the influence of ENSO on the climate of the altiplano appears to be considerably more complex and uncertain than in coastal regions in the North.

Forecasting El Niño

The semiperiodic occurrence of El Niño and La Niña events and their important influence on the climate of the region has spurred interest in the applications of seasonal forecasts.

In their forecasts, scientists focus on particular regions of the Pacific that have shown a strong correlation with the evolution of an El Niño (and La Niña) event. For Peru, two regions are of particular interest: the coastal Pacific and the equatorial Pacific.

As highlighted earlier, El Niño and La Niña are now recognized as components of


the El Niño Southern Oscillation (ENSO), an ocean atmosphere phenomenon, which spans the entire Pacific basin and has global effects on the climate. The evolution of global El Niño and La Niña events is closely associated with SST anomalies along the equatorial Pacific (as well as changes in the

atmospheric pressure differences between the eastern and western Pacific regions).

International forecasts of global El Niño events predominantly concentrate on regions in the equatorial Pacific, which have shown high predicative capacity, such as the NINO 3, 4 and 3.4 regions (Figure 4). Climate models driven by observed SSTs can now be used to forecast, with considerable skill, the evolution of El Niño events more than six months in advance (Figure 5).

Internationally, institutes such as the International Research Institute for Climate and Society (IRI) use SST thresholds that results in a 25%-50%-25% distribution of La Niña, neutral, and El Niño


Figure 4. The NINO Regions



Note: Temperature anomalies observed in the NINO 3.4 and NINO 3 and 4 regions are predominantly used in forecasting ENSO events. The NINO 1+2 region plays an important role in influencing the climate of northwestern Peru, but is less well suited for seasonal climate forecasts.

Source: IRI, <http://iri.columbia.edu/climate/ENSO/background/monitoring.html#Niño>.

Figure 5. NINO 3.4 Forecasts: Lead Time by Month and Predictive Skill



Note: Depending on the month of observation, forecasts with a high predictive skill can be made of more than six months in advance of the event.
Source: IRI, Columbia University.

conditions over a base period from 1950–2002. This distribution reflects the number of global ENSO events that have historically been recognized by experts. In order to arrive at such a distribution, the SST thresholds have to be adjusted throughout the year to account for seasonal fluctuations in temperature (Table 3). Based on these thresholds, seasonal forecasts are developed. While this classification system helps to identify the years that have commonly been

accepted as global El Niño or La Niña years, they may not necessarily always trigger climatic changes in northwestern Peru that matter to local livelihoods.

NOAA's climate prediction center uses a slightly different system for classifying warm (El Niño) and cold (La Niña) episodes of ENSO, which are based on thresholds above or below $\pm 5^{\circ}\text{C}$ average seasonal SST sustained for a minimum of three consecutive months in the NINO 3.4 region, respectively. This results in a similar historical distribution of El Niño, normal, and La Niña years (see Appendix B).

The main influence on interannual variability in northwestern Peru is associated with the NIÑO 1.2 region (Figure 4). The predictive skill for the NIÑO 1.2 region, however, is lower and, hence, forecasts have shorter lead times. Consequently, the SST thresholds often used for determining the evolution of a global El Niño do not necessarily always translate into a strong local El Niño in Northern Peru, which is associated with torrential rains and flooding conditions.

While changes in the NIÑO 3.4 region are certainly important to the climate in Peru, it is important to recognize that other thresholds are better suited for El Niño conditions in Northwestern Peru and coastal Ecuador (Pilar, personal communication, Jaimez, personal communication). Local and regional scientists certainly recognize this problem. Usually a combination of indicators and different thresholds are used by the national and regional centers when issuing

Table 3. Three-Month Temperature Anomalies for Classification of El Niño (EN) and La Niña (LN) Conditions, in Degrees Celsius

Season	LN	EN	Season	LN	EN
DJF	-0.65	0.65	JJA	-0.50	0.45
JFM	-0.55	0.50	JAS	-0.50	0.45
FMA	-0.45	0.40	ASO	-0.55	0.50
MAM	-0.40	0.40	SON	-0.75	0.70
AMJ	-0.45	0.45	OND	-0.75	0.75
MJJ	-0.50	0.45	NDJ	-0.70	0.75

Source: IRI, Columbia University.

their forecasts; however, this information is often not published in international scientific journals.

The altiplano of Peru is predominantly influenced by changes in the equatorial in the SST along the equatorial Pacific; i.e., changes in the NINO 3.4 region.

Despite the widespread influence of ENSO on climatic conditions, there is no uniform definition of El Niño conditions that automatically would apply also to local climatic conditions. In the case of Peru, the experience developed by national and regional meteorological organizations in identifying thresholds that are locally relevant is of considerable importance. SENAMHI and other organizations do evaluate the accuracy of forecasts; however, much of this knowledge and information on thresholds is not published internationally or is at least publicly available.

As there are different classifications for El Niño events based on the ever-evolving indexes and practices on the national and international level, this may lead to conflicting messages. The Climate Outlook Forums represented a promising effort to develop a consensus view on the status of El Niño events in various regions of the world (IRI 2001); however, it has not translated into continuous coordination efforts for scientific forecasts in this region.

Given the strong influence of ENSO on climatic characteristics, forecasts have not only focused on alerting for El Niño events, but also on providing more continuous insights into the climatic conditions of the region. These forecasts are presented as probabilistic assessments of whether a particular region will be faced with drier, wetter, or average climatic conditions.

Institutional Structures for Disaster and Climate Risk Management

Over the last five years, there has been an increasing shift away from a reactive approach predominantly focused on disaster mitigation and toward building

an integrative framework that also strongly focuses on disaster prevention, by identifying risks and vulnerabilities (World Bank 2006).

The Plan Nacional de Prevencion y Atencion de Desastres (National Plan for the Prevention and Attention to Disasters) was approved in 2004. The plan includes objectives and identifies specific strategies, aimed at improving the risk assessment, vulnerability reduction, integration of risk management into public planning, strengthening institutions and the participation of communities. However, there are currently no clear definitions of the responsibilities of the individual institutions and their specific operational roles (World Bank 2006).

Given the vulnerability to hydrometeorological hazards and the strong influence of ENSO, the growing emphasis on vulnerability reduction is also reflected in institutional efforts to collect and disseminate climate information, aimed at reducing the vulnerabilities of climate-sensitive sectors.

Disaster preparedness efforts focused on ENSO events are coordinated by the National Study of the El Niño Phenomenon (Estudio Nacional del Fenomeno El Niño; ENFEN), which addresses the national component of the Regional Study of the El Niño Phenomenon (Estudio Regional del Fenomeno El Niño; ERFEN). ERFEN was established as a program by the Permanent Commission for the South Pacific (Comisión Permanente del Pacifico Sur) in 1974 and guided by national committees from its four member countries. Aside from Peru, these include Colombia, Ecuador, and Chile (Garcia-Godos et al. 2004).

ENFEN Peru is comprised of six national institutions (Table 4). The information flow between the various institutions is presented in Figure 6.

The climate and oceanic conditions are monitored by IMARPE, IGP, DHN, and SENAMHI. Based on a collective and consensus-finding process, El Niño forecasts are being issued. Smaller-scale interventions

Table 4. Members of ENFEN–Peru

<i>ENFEN Institutions</i>	<i>Focus</i>
IMARPE: Instituto del Mar del Peru	Climate Monitoring
IGP: Instituto Geofísico del Peru	
DHN: Direccion de Hidrografia y Navegacion (within the Navy)	
SENAMHI: Servicio Nacional de Meteorologia e Hidrologia	
INDECI: Instituto Nacional de Defensa Civil	Disaster Prevention & Mitigation
INRENA: Instituto Nacional de Recursos Naturales	

Source: J. Garcia-Godos and others, 2004. Prepared as part of a project initiative.


are often implemented by members of ENFEN itself, while larger-scale projects are usually presented by INDECI to the relevant ministries, such as the Ministry of Transport (Garcia-Godos et al. 2004).

INDECI and INRENA both work on local levels (Garcia-Godos et al. 2004). This provides the opportunity to engage in awareness raising, feedback, and information exchange with local populations

through the existing local institutional networks of these executive agencies.

The institutions most directly engaged in the dissemination of climate information are SENAMHI, INDECI, and INRENA. SENAMHI, as the national meteorological service, monitors climate and weather conditions for the entire country. The General Directorate for Agrometeorology (GDA) situated

Figure 6. Diagram of Information Flow through ERFEN Network in Peru



Source: J. Garcia-Godos et al. 2004, prepared as part of this study.

within SENAMHI, focuses on the generation of agrometeorological forecasts. Relevant data on crop productivity is collected on a weekly basis by a locally recruited agent, who is usually a local farmer trained by SENAMHI. The data is then usually transmitted via phone, which also allows for passing on other potentially important information, such as local knowledge of environmental indicators. Consultations with GDA suggest, however, while staff are aware of this knowledge, there are currently no systematic efforts to collect and evaluate such knowledge and compare it with scientific forecasts (Garcia-Godos et al. 2004).

GDA also monitors the phenological development of crops in relation to weather conditions in more than 400 observatories across the country. The information gathered by GDA is made publicly available in three types of bulletins: one bulletin focused on short-term trends is published every 10 days, and two monthly reports, focused on hydrometeorological and agroclimatic conditions, also provide information on tendencies and forecasts.

While SENAMHI aims to provide information for authorities and farmers, the main user of the climate information collected by GDA is indeed the Ministry of Agriculture (MINAG), and the General Directorate of Agrarian Information edits the information into a more user-friendly format and distributes it to its regional directorates. The detailed agroclimatic information, however, requires access to the internet. A pilot project has been launched by SENAMHI and MINAG in the Cañete Valley of Peru to assist farmers with Internet access to utilize this information for improving crop production. In general, the distribution channels of these forecasts are still limited, as this requires adequate literacy and access to a computer (Garcia-Godos et al. 2004). Consequently, the main communication of climate information occurs through radio broadcasts, which receive information updates from SENAMHI.

MINAG combines statistical production data with climate information from SENAMHI. The agricultural data is collected by staff from agrarian agencies working at the district level, which constitutes the lowest administrative unit in Peru. The data is based largely on assessments and estimates and does not include systematic surveys (Garcia-Godos et al. 2004). After the analysis, the information generated by MINAG is then passed on to agrarian agencies at the provincial and district level. The information, however, does not always get there due to bottlenecks in the distribution path. Other information arrives with a time lag of 20 days or more (Garcia-Godos et al. 2004). Consequently, the forecasting aspect for the end user may be lost, limiting the practical relevance of the information in guiding strategies aimed at vulnerability reduction or disaster preparedness.

Climate Change — Challenge and Institutional Response

Over the past 100 years (1906–2005), the earth's annual surface temperature has increased by about 0.74°C (IPCC 2007). The rate of warming is accelerating. A total of 11 of the 12 warmest years on record have occurred in the past 12 years, and the projected increase in global mean surface temperature between 1990 and 2100 is projected to lie between 1.0 to 6.3°C. As global warming of this century will be more pronounced than the trends observed in the last century, it is expected to bring more drastic changes in climate and associated environmental responses.

Given the diverse climates and ecosystems, steep topographies, existing disaster risk, and high dependency of livelihoods on natural resources in Peru, climate change poses an additional threat to the sustainable development prospects of the country. There is a need to understand how climate change will impact climatic conditions across temporal and spatial scales, affect the provision of ecosystem services, and alter the exposure to climatic extremes. Given the influence of ENSO on Peru's climate and disaster risk,

it is furthermore important to gain insights into the implications of global warming on the occurrence and magnitude of El Niño and La Niña events. Improving such knowledge will be essential to guiding longer-term development-planning processes and complementing existing efforts aimed at reducing vulnerabilities to current climatic risks.

It is interesting that Peru's efforts in climate change so far have been more focused on the international policy dialogue and on mitigation options for the emission of greenhouse gases (World Bank 2006). Only recently has Peru increased its emphasis on identifying and implementing adaptation measures to climate change.

The institutional landscape for managing environmental and natural hazards in Peru is undergoing change itself. The country is setting up for the first time a Ministry of Environment. Most of the efforts to mainstream climate change policies into national development in Peru have been led by the Consejo Nacional Ambiental (National Council for the Environment (CONAM), which has been chairing the National Commission on Climate Change. The new ministry will place a strong emphasis on addressing climate change, likely building CONAM's experiences through the PROCLIM project and emphasis of a holistic approach, which focuses on mitigation (reduction of greenhouse gases) and adaptation.

As part of its efforts to assess climate change risks, CONAM has also assessed the potential implications of climate change for the occurrence of El Niño events. Using a range of climate change models, CONAM found an increase of the SST in the NINO 3 region under global warming scenarios (Figure 7), which suggests that conditions may become more conducive to the evolution of El Niño events.

The World Bank is engaged in research focused on identifying climate change impacts and vulnerabilities and developing project activities that promote adaptation to climate change in Peru and other Latin

American countries (see Vergara 2005 for details on project activities).

Of major concern are the implications of climate change for the water balance for the Andean region and the associated economic consequences (Vergara et al. 2007). The majority of Peru's population and economic activities are located along the coast and in highlands. Both regions are predominantly dry and highly dependent on runoff from the cordilleras. Recent research confirms a massive retreat of glaciers, which is in line with overall warming trends and an upward shift in freezing point isotherm (Kaser 2001, Francou et al. 2003). The retreat of glaciers results in an initial temporary increase in runoff, followed by a decline and shutoff of water supply as the glacier disappears. Given that glaciers have been shown to contribute considerably to the overall water balance in river basins in the Andes, the downstream implications of a decline in this freshwater resource represents a considerable and growing challenge to agricultural activities and hydropower generation in the region (Vergara et al. 2007).

Climate Change and El Niño


Another concern is the influence of climate change on atmospheric and oceanic circulation patterns and associated consequences for such phenomena as ENSO. Current scientific thinking suggests that the climate will enter a weak El Niño-like mean state, with El Niño events resembling strong El Niño events of today (e.g. IPCC 2001a). This is echoed by results of a recent climate change study conducted by CONAM, which shows a warming trend in SST for NINO 3, a region where SST anomalies are correlated with the occurrence of ENSO events (Figure 7).

Since 1976, more warm El Niño episodes than cold La Niña episodes have been observed, but it is currently not clear whether this can be attributed to climate change.

With more El Niño–like general climatic conditions and an increased frequency of strong El Niño events, one may expect wetter conditions in Northern Peru and

drier conditions in the altiplano, but the link between El Niño and climate change is not fully resolved and hence such effects are still largely speculative.

Figure 7. NINO 3 Region Climate Trends



Note: Climate simulations suggest a general warming trend for the NINO 3 region. As positive anomalies in SSTs are associated with El Niño conditions, this may suggest a shift toward a more El Niño–like climate conditions. The colors show different simulation runs by Global Climate Models for the NINO 3 region, all of which point toward an increase in sea surface temperature. *Source:* CONAM.

2 Climate Resilient Development— Community Perspectives

The previous sections have presented an overview of the exposure of Peru to natural hazards with a strong emphasis on the influence of El Niño on climate risks and then described the existing institutional structures for disaster prevention and preparedness. In this context, climate change has been recognized for its threat to sustainable development due to its potential implications for the natural and water resources effects and the associated consequences for Peru's economy.

In light of climate change, the existing vulnerabilities associated with the annual and interannual climate variability and ENSO underscore the importance of adaptive measures that enhance the resilience of livelihoods and economic processes in Peru. Adaptation to climate change hence has to begin with better managing current climate risks and should be viewed as a continuous process. Adaptive measures range from strategies, policies, and investment frameworks enacted at the national level to efforts at the local level. The time horizon of adaptive strategies and measures has to be tailored according to the scope of the activity and capacity of the target groups.

This chapter focuses on how communities perceive the risks of their environment and how they cope and adapt. The experiences of communities in living in the department of Piura, northwestern Peru, are juxtaposed with the situation of communities in the department of Puno, representing the Peruvian altiplano. Both regions have in common an interannual climate variability influenced by ENSO. The effects of El Niño and La Niña on the climate, however, differ considerably, as

does the overall risk profile. Based on these responses and in recognition of national risk management structures, adaptive capacity-building needs aimed at increasing the resilience to climate variability and change are discussed.


Approach

Under coordination by the Centro Internacional de la Papa (CIP), the community assessments were carried out by the Intermediate Technology Development Group (ITDG) in the department of Piura and Centro de Investigación de Recursos Naturales y Medio Ambiente (Research Center for Natural Resources and Environment; CIRNMA) in the department of Puno.

In Piura, a total of four communities were selected covering a transect from the coast to the sierras in the proximity of the Piura river (Map 2, Piura). The assessment included the coastal community of Parachique (sea level, bajo Piura), Chato Grande (30 m a.s.l, bajo Piura), Locuto (80 m a.s.l, medio Piura) and Cabuyal in sierras of Piura (1,470 m a.s.l., alto Piura). A description of the individual community characteristics is provided in Table 5.

For the assessments in Puno, six communities in different proximity to Lake Titicaca were selected, including the communities of Santa Maria (3,862 m a.s.l), Ancacca (3,950–4,050 m a.s.l.), Yanamocco (3,841 m a.s.l.), Candile (3,834 m a.s.l.), Cari Cari (3,950 m a.s.l) and Alto Achuyo (3,947 m a.s.l.). Further information on the individual communities is provided in Table 6, and spatial information

Map 2. Location of Communities Assessed in Piura



concerning the location of communities relative to Lake Titicaca is given in Map 3.

A participatory process was chosen to elicit information from the communities. The assessments first focused on the perceptions of each community concerning their exposure to environmental risks and hazards. Participants were asked to recall events that constituted a major disaster to the community by significantly affecting human health, infrastructure, or production systems.

Table 5. Community Characteristics, Piura

Department of Piura, N-Peru


Parachique (5°45'51" S/ 80°51'53" W, sea level): Representing bajo (lower) Piura, this coastal community located at the fringes of the Sechura desert largely depends on fishing activities. Four types of fishermen can be distinguished: artisan fisherman, who use lines or nets; shellfish divers; boat owners; and industrial fishermen, who harvest fish of up to 50 metric tons (MT), predominantly for the fish meal industry. The poorest segments of Parachique's inhabitants do not participate in fishing activities and solely depend on dryland farming and products of the dry forest. Many women obtain income by processing shellfish.

Chato Grande (5°21'20" S/ 80°40'3" W/ 30 m a.s.l.): An inland community of bajo Piura, Chato Grande is located in the lower watershed of the Piura river. Fertile soils are interspersed with tropical dry forest. Having access to irrigation, the community's main activity is crops (cotton, rice, maize, beans). Goats and poultry are also raised.

Locuto (4°54'53" S, 80°20'56" W, 80 m a.s.l.): Located in the tropical dry forest in medio (middle) Piura, the carob tree (algarrobo, *Prosopis limensis*) plays a central role in the livelihood of Locuto's inhabitants, who harvest the beans and also use the tree for fuelwood and charcoal. The carob tree is also an important resource for handicrafts (along with two other tree species: *Cordia rotundifolia*, *Cappais angulata*). Farming activities are concentrated in the riverbed areas, which have sufficient moisture for planting sweet potatoes, maize, beans, and various fruits. Depending on rainfall, planting activities (mainly beans and watermelon) are occasionally extended in forest areas cleared for production. Livestock includes predominantly sheep and goats. Beekeeping is also common activity.

Cabuyal (5°02'17" S, 79°49'42" W, 1,470 m a.s.l.): The area of this highland community of alto (higher) Piura extends from 1,400–2,000 m above sea level, with nucleated population center being located at around 1,470 m a.s.l. Agricultural activities are diversified, including dryland farming of maize, wheat, beans, and green peas, along with fruit trees and coffee at lower elevations. Livestock production is also of importance. With respect to climate, it is important to note the specific characteristics of this part of the sierras. At these lower elevations of the sierra, southeast winds bringing humidity from the Amazon meet westerly airflows from the Pacific. As a result, this area experiences heavy rainfalls, which is unique in the highlands.

In Piura, the community members voted on the

Map 3. Location of Communities in Puno

severity of the events and ranked them accordingly. In Puno, the communities first ranked the hazards according to their level of concern and general impact. As the number of hazard events recalled by the communities in Puno was too high, it was decided to solely focus on major disaster events. A time line for these events was then constructed. The individual events were discussed in focus groups, including stakeholders involved in managing risks or who most affected by the event. In Puno, CIRNMA also developed spatial risk maps with the focus groups, detailing the impact zone of major disaster events in the communities.

Following the discussion of hazard risks, disaster events, and vulnerabilities, the discussions then concentrated on the availability and use of weather and climate information. Community members were asked whether and how they obtain access to external information, the trust they place in the various information sources, and whether and how they respond to the information content. In this context, community participants were

also presented with various information formats and asked to provide feedback on the value of this information for practical information needs.

Besides formal information sources, community representatives were asked to describe the type of local indicators or other informal sources of information that they consider of value in warning of climatic extremes or predicting the climate characteristics of forthcoming seasons.

The assessments then discussed which coping and adaptation mechanisms the community members currently employ to lessen their vulnerability to and the impact of extreme climatic events.

The goal of the discussions was also to identify the current challenges and barriers communities perceive in managing climatic conditions. These discussions focused on information needs, constraints within the community, and external barriers.

Based on these discussions, each community then developed recommendations of action that could be undertaken without and with external assistance to strengthen the resilience to climatic hazards. The results of the individual community assessments were then presented in regional workshops in the cities of Piura and Lima, where community representatives from Piura and Puno were brought together with representatives from governmental institutions and researchers engaged in disaster preparedness. Together the participants identified cross-cutting issues and prioritized the core issues that needed to be addressed.

Table 6. Community Characteristics, Puno

Department of Puno, altiplano, S-Peru

Santa Maria (16°07'08" S, 69°39'30" W, 3,862 m a.s.l.): This Aymara peasant community is located 4 km away from Ilave, consisting of 130 families. In recent years, young adults are migrating to Tacna, Ilo, and Arequipa in search of work. The community is mainly dedicated to crop production. Approximately 70% of the land is planted with potato, quinoa, ollucos, and forages like alfalfa and oats. Land constitutes a limiting resource, with the community covering 340 hectares (ha). Production areas include pampa (60%), hillside (20%), and hills (20%). Approximately 25 ha are cultivated. Cattle, sheep, and small animals represent complementary income sources. On average, a family owns three cattle, six sheep, two pigs, four chicken. Some families also have guinea pigs.

Terraces are used or being built for agricultural activities. There is no access to irrigation. Planting decisions are made through the aynoka system, meaning decisions on when and where to plant are made collectively, but the farming activity is made collectively. Organizations working in the community: PRONAMACHS, mother's club.

Ancacca (16°07'42" S, 70°02'22" W, 3,950–4,050 m a.s.l.): The peasant community of Ancacca has a population of 66 Aymara families, and is 38 km southwest of the city of Puno. It belongs to the Pichacani district, in the province of Laraqueñi. It has 842.5 ha of land of individual access, and 270 ha given through agrarian reform for communal use. It has a board that oversees the community. Two-thirds of the land is managed individually, while a third is communal. Even though some crops are grown in this region, the risk of loss is high, making livestock production their main economic activity. Seventy percent of the land is in the pampas. There is cultivation of potato, quinoa, and cafiwa for food, and forages for livestock. Wells for irrigation were constructed in 1987, but these do not work well. The community has a communal building, a preschool, an elementary school, a storage facility for potato seed, and dip for sheep. There is electricity in the homes, and a reservoir of potable water. The main institution working with the community is the Corredor Económico Puno-Cusco (Economic Corridor of Puno-Cusco) which provides technical assistance for livestock, handicrafts, and husbandry of small animals. This community has a strong organization, meeting every 30 days. In the last year, they suffered from frost and drought from January to March (growing season). Young people migrate to other provinces in search of farming jobs.

Yanamocco (5°40'28" S, 70°05'15" W, 3,841 m a.s.l.): This peasant community is located in the Suni A agroecological zone, 35 km from Lake Titicaca in the province of Atuncolla. It is a Quechua community of 43 families that are registered, and 44 that are not. This community has approximately 133 ha. It also has land received during the Land Reform. It has 470 ha of grazing land appropriate for South American camelids, and 96 ha of agricultural land. Most of Yanamocco is located in the pampas, in the watershed of river Illpa. The land is suitable for pastures such as alfalfa. The limited crop production is mainly for household consumption. It has its own governing committee that includes defense, rondas campesinas (peasant patrols), an el vaso de leche committee (glass of milk committee), a committee for electricity, one for forages, and a mother's club and neighbor's committee. Because most of the landscape is pampa, it is prone to floods and frost. Dairy production is the main economic activity. CIRNMA, CARITAS Puno, and the municipality have been working on forage production. There is also processing of dairy products and an improvement in cattle breeding. All of the production is in the pampa. The community has some raised fields (Waru Warus), a pre-Inca technology, to manage flood risks and improve micro-climatic conditions for crop planning, and eight terraces (Andenes).

A study of economic conditions (CIRNMA 2004) showed that families with greater resources have on average 25 cattle, 62 sheep, 9 pigs, 16 chicken, and two to 10 alpacas. Those in the middle of the economic spectrum have 13 cattle, 22 sheep, three pigs, and three chickens, and some have alpacas. Those with lesser resources have on average seven cattle, 11 sheep, one pig, seven chickens, and no alpacas. Regarding infrastructure, they own a local community building, a potato seed center, a health facility, and a preschool facility (daycare, or wawahuasi). Organizations in the region work with CARE on training in agriculture, CIRNMA on credit and training in livestock production, the Corredor Económico Puno-Cusco on training and technical assistance, Caritas Puno on setting up cultivated pastures, and the Ministry of Agriculture on training in animal production and health. Some families are producing yogurt and cheese. The resource-endowed women work on handicrafts such as blankets, sweaters, and socks in the evenings.

Candile (15°35'31" S, 69°56'47" W, 3,834 m a.s.l.): Candile represents a sector of the larger peasant community of Carata. The population is predominantly Quechua and consists of 100 families and an additional 10 migrant families. Crops are the main activity. These include potato, quinoa, cafiwa, barley grain and forage, and oats. Livestock production also plays an important role. There are approximately 740 cattle, 900 sheep, and 150 pigs. Fishing represents an additional income source. The total extension of land is 610 ha, of which 300 ha are used for grazing and 300 ha for crop planting. Waru warus, or raised fields, a pre-Inca technology to manage water floods on crops fields, are found in Candile (in the community of Yanamocco).

Cari Cari (15°46'33" S, 70°21'42" W, 3,950 m a.s.l.): The population center of Accopata, Cari Cari, is a Quechua community located 40 km northeast of Puno. This is not a registered center. It belongs to a peasant organization, Central Accopata, recognized in 1967. It is located in Mañazo, 40 km northeast of Puno. It has approximately 100 families. Cari Cari has 212 ha managed individually, and 1,425 ha that are managed by the community. It has mostly a pampas landscape. There are 160 ha suitable for crops. Potatoes, quinoa, and faba beans, along with forages, are the main crops. Cattle, sheep, and alpaca are the main types of animals kept. Wealthier families own on average 30 cattle, 55 sheep, six alpacas, and two pigs. A middle-income family owns nine cattle, 17 sheep, three alpacas, and one pig. Low-income families own three cattle, 14 sheep, two alpacas, and one pig, on average. Young adults migrate to other regions when natural disasters occur. Only a few families own trees. There is a primary and a secondary school, although the latter is too small to adequately accommodate all children. There is a preschool and a health building, but no health care provider. The community is currently working with CIRNMA on microfinance and livestock technical support, and with the Universidad Nacional del Altiplano Puno on potato production al partir (sharecropping). Caritas Puno and the municipality of Mañazo have been working there since 2002 to establish forages). PECSA provides support on animal health. Youth migrate in search of work when there are unfavorable climatic conditions for agricultural activities.

Alto Achuyo (16°14'28" S, 69°01'45" W, 3,947 m a.s.l.): Representing an Aymara population center, 100 families live in this community, which extends across 210 ha of land. Alto Achuyo represents one of six sectors of Achuyo, which is recognized as a community and has its own public officials. The hillsides of the community are predominantly used for cultivation. Alto Achuyo is located near Lake Titicaca and the Winaymarka. Both lakes influence the climate of the community. Crops constitute the main activity, followed by livestock and handicrafts. The crop rotation consists of potatoes, followed by oca, faba beans, wheat or barley, forages, Tarwi, and fallow. All families own plots on all of the hillsides in the aynocas. The community has old terraces that are still used. No irrigation projects exist. Families usually possess three cattle, 15 sheep, four pigs, and three chickens, on average. Organizations working in the community include PRONAMACHS, Intervida, the church, and PRADERA.

Perspectives from Communities in Piura and Puno

The following sections are structured around the following themes: 1) Disaster Risks and Impacts, 2) Vulnerability, 3) Coping with and Adapting to Climate Hazards, 4) Information: Weather Forecasts, Early Warning, and Seasonal Climate Outlooks, and 5) Traditional Information and Knowledge. Each theme is briefly introduced in its relevance to climate resilient development. The results of community assessments in northwestern Peru and Piura are then presented and subsequently discussed, drawing on data records, publications, and complementary interviews.

Disaster Risks and Impacts

This section sets the stage for the subsequent discussions by outlining the exposure to natural hazards and impacts on the livelihoods of the community. The community assessments outline their personal perceptions of disaster risks and how the individual hazard types have translated into disaster events that have adversely affected human health, assets, or development prospects. The disaster impacts of the communities are not only a function of the exposure to a particular hazard event, but also the outcome of the underlying vulnerability factors. Therefore, the discussions here are closely linked to the subsequent sections on vulnerability, coping and adaptive capacities, and information access.

Piura

The recollection of disaster events by the communities in Piura illustrate the challenge of coping with too wet and too dry conditions (Table 7) in the coastal (bajo Piura) and lowlands (medio Piura). Cabuyal, the community assessed in the sierra (alto Piura), is exposed to extremely wet conditions. It suffers less from too dry conditions as the rainy season is more consistent, but the community representatives highlight other climatic extremes that frequently destroy their assets.

Concerning the relevance of El Niño in the risk perceptions of the communities, it is striking that only strong El Niño events are recalled as disasters. The communities either directly identified these events as disasters (Parachique) or clearly linked the observed heavy rains and flooding to El Niño in the recollection of disaster events (Chato Grande, Locuto, and Cabuyal).

Cabuyal is the only community that recalled a disastrous rainfall episode that was not linked to El Niño. The heavy precipitation experience in 2001 was in fact considered the most severe disaster in the recollection of the community.

The flooding event experienced by Chato Grande in 2002 was not linked by the community to El Niño (despite being classified as an El Niño year; see Appendix B). The community associated El Niño and associated floods with intense rainfall episodes, which did not occur as that time. 2002 was the first time in the memory of the community participants that a flood occurred without preceding intense rainfall.

The communities all highlighted various drought episodes that significantly impacted their livelihood. During the time of the assessment (2005), the current drought period was a shared concern. The starting and end dates of the dry periods assigned to the recent and preceding drought periods differed between the communities. Aside from the recent dry period, the 1960s, and in particular, 1968, were remembered as times when the community severely suffered under drought conditions, as highlighted in the assessments in Chato Grande and Cabuyal. Other dry periods were individually reported by the communities, but not shared. The coastal community of Parachique also equates dry periods with colder-than-usual temperatures.

In Cabuyal in the sierras of Piura (alto Piura), experiences also highlighted a range of other climatic hazards, which are experienced on an annual basis

and hence constitute a consistent challenge for the community. These include annually recurring fogs during the rainy season and strong winds. Nonclimatic events, which were recorded as disasters in the discussion, include a forest fire (Locuto), and earthquakes (Parachique, Locuto, and Cabuyal), as well as cholera epidemics (Parachique, Locuto).

The responses from all four communities show a persistent vulnerability to climatic hazards. Associated with the range of predominantly climate-related hazards experienced by communities, impacts on infrastructure, human health, education, and economic activities are recurrent themes (Table 8).

Of the major disaster events recalled by the communities, years with strong El Niño events are featured in the ranking of all communities. There is a clear understanding that El Niño is associated with excessive and prolonged rainfall episodes, but

it is predominantly the strong events of 1982–83 and 1997–98, where the negative effects outweighed the positive impacts. During these events, the initial destructive impacts associated with the floods and heavy rainfalls predominate. More moderate events in the 15-year period between these two strong El Niño events were not highlighted as disasters in the discussion. In fact, according to the perception of communities, if the households are able to cope with the floods and heavy rain, the subsequent effects of El Niño are desirable, as they represent the opportunity to increase agricultural and livestock production.

Prolonged dry conditions are generally perceived as negative. The only exception in the assessments is the fishing community in Parachique, where most livelihoods benefit from favorable fishing conditions. None of the communities, however, relate dry conditions to La Niña events. There appears to be

Table 7. Disaster Events, as Recalled and Ranked by Communities in Piura, Peru

Ranking	Community			
	Parachique	Chato Grande	Locuto	Cabuyal
1	Floods, climatic extremes (1982–83 El Niño)	Floods (1997–98 El Niño)	Dry spells & droughts (2001–05)	Heavy rains (2001)
2	Dry spells, droughts (1998–2005)	Floods (1982–83 El Niño)	Flooding and climatic extremes (1997–98 El Niño)	Drought period, below normal rains (2003–05)
3	Cholera, 1992	Piura river flooding (2002, not recognized as an El Niño year by community)	Forest fire (1998)	Strong winds (annual problem)
4	Floods, climatic extremes (1972 El Niño)	Drought periods: 1960–68, 1972–80, 2003–05 (equally ranked by community)	Flooding and climatic extremes (1982–83 El Niño)	Fog (annual problem impacting crops)
5	Dry spells, droughts (1970–85)		Cholera (1992–93)	Floods, heavy and prolonged rains (1997–98 El Niño)
6			Earthquake (1972)	Floods, heavy and prolonged rains (1982–83 El Niño)
7			Earthquake (1970)	Earthquake 1970
8				Drought (1968)

no clear conceptual understanding of La Niña as the counterpart of El Niño.

Table 8. Impacts of Climate Hazards Identified in the Community Assessments in Piura

<i>Event</i>	<i>Impacts Highlighted in:</i>			
	<i>Parachique (coastal, bajo Piura)</i>	<i>Chato Grande (bajo Piura)</i>	<i>Locuto (medio Piura)</i>	<i>Cabuyal (alto Piura)</i>
El Niño	<p>Housing & infrastructure:</p> <ul style="list-style-type: none"> Destroyed houses (eastern sector of bay most affected) School building destroyed Isolation (road to Sechura and bridge affected) <p>Education:</p> <p>Delay in education (classes suspended in 1972 & 1983)</p> <p>Health:</p> <ul style="list-style-type: none"> Increase in vector-borne diseases Malnutrition; food supply low during flooding events <p>Fisheries:</p> <p>Negative impact for small-scale fisheries, which suffer from migration of fish stocks</p> <p>Agriculture:</p> <p>Initial adverse impacts from floods are followed by subsequent benefits from revitalized dry forest and ecosystems. Increased pasture and planting areas.</p>	<p>Housing & infrastructure:</p> <ul style="list-style-type: none"> Collapsed houses; impact on school building After flooding in 1983, 24% of the community was relocated to Nuevo Chato Grande, which is perceived as a disintegration of the village's social structure Loss of schooling time due to building damage isolation. Children were withdrawn from classes for extended period due to the adverse economic situation of some families. <p>Health:</p> <p>Increase in vector-borne diseases, lack of clean water associated with increase in diarrhea, and respiratory problems.</p> <p>Agriculture & livestock:</p> <ul style="list-style-type: none"> Loss of crops and seeds, destroyed harvest of the long (Feb & March) and short (July) planting season in 1997–98; similar impact in 1982–83. During heavy rains and floods, loss of pasture. Increase in pests and animal diseases; decrease in livestock Irrigation channels were partially destroyed or silted up. Migration of resource-poor households Longer-term positive effect: Rains reactivated occasional fields, where farming is dependent on rain. Increase in pasture after floods receded; increase in livestock 	<p>Housing & infrastructure:</p> <p>Isolation</p> <p>Health:</p> <p>Increase in malaria</p> <p>Agriculture & livestock:</p> <ul style="list-style-type: none"> Loss of crops; during the 1997–98 event, the Piura river changed course and destroyed fields in its path; compounding effects of increases in pests. Afterward, agrochemicals were introduced to improve yields. High mortality in livestock due to lack of protection. Compounding effects from increases in disease and parasites. <p>Positive effect:</p> <ul style="list-style-type: none"> Revitalization of dry forest Reactivation and expansion of occasional fields. <p>El Niño events are generally recognized for their positive longer-term effect on agriculture and livestock activities.</p>	<p>While Cabuyal usually receives considerable rainfall during the rainy season, El Niño events increase the length, and sometimes also the intensity, of the rainy season, and the preceding months are characterized by high levels of humidity (Angulo, personal communication)</p> <p>Infrastructure & housing:</p> <p>Isolation due to collapse of Carrasquillo bridge and road blockage from landslides</p> <p>Agriculture & livestock:</p> <ul style="list-style-type: none"> Loss of crops and seeds Loss of fertile land (landslides and erosion) Increases in fungal diseases; about half of the crops were lost in 1997–98 Loss of livestock Increase in livestock diseases (e.g., anthrax and foot and mouth, and emergence of others, such as Tupe) Decline in livestock sale price Increase in cattle theft and crime <p>Community structure:</p> <p>Migration of resource-poor household members in search for additional income</p> <p>Positive, subsequent effect:</p> <ul style="list-style-type: none"> Increase in pastures Increase in livestock numbers

<i>Event</i>	<i>Impacts Highlighted in:</i>			
	<i>Parachique (coastal, bajo Piura)</i>	<i>Chato Grande (bajo Piura)</i>	<i>Locuto (medio Piura)</i>	<i>Cabuyal (alto Piura)</i>
Drought	<p>Agriculture & livestock: No rain-fed agriculture possible; degradation of dry forest; decimation of livestock</p> <p>Beneficiaries: The cold and nutrient-rich waters improve fishing conditions. Families can increase income.</p>	<p>Agriculture & livestock:</p> <ul style="list-style-type: none"> • Impact reduced due to access to irrigation. • Planting and crop selection constrained. • Lack of pasture • Drop in milk production, increase of animal diseases and mortality • Drop in sale prices for livestock <p>Education: Resource-poor families withdraw children from school to help with income-generating activities (male children over eight, girls did not receive education because of this situation)</p> <p>Community structure: Resource-poor families or families who lost land during preceding 2002 flood could not generate enough income; family members forced to temporarily migrate.</p>	<p>Human health:</p> <ul style="list-style-type: none"> • Malnutrition, especially in children, associated with a drop in school performance. • Increase in gastrointestinal and respiratory diseases associated with lack of water and bad quality. <p>Agriculture & livestock:</p> <ul style="list-style-type: none"> • Lack of water made it impossible to plant in the floodplain (usual planting area) in 2002 and 2003; in 2004, these fields could only be planted using a reduced sowing rate • Scarcity of pasture and food for animals. • Lack of water • Increases of diseases such as plague and diarrhea and parasites in animals; malnutrition of livestock, loss of weight • No milk production from goats • Drop of sales price, buyers exploit situation • Quantity of syrup from the carob tree in the dry forest reduced; reduced production of honey. <p>Environmental Degradation: Increase in deforestation due to the collection of fuelwood as alternative income source.</p>	<p>The drought of 1968 is present in the collective memory of the community, as the absence of rain triggered widespread migration. The recent dry period is associated with below-normal rainfall conditions for several years.</p> <p>Agriculture & livestock: Pastures dried up in 1968; farmers were forced to sell livestock or land to obtain food and compensate for losses. No milk and cheese production. The recent dry period is less severe but has led to a significant decline in agriculture output.</p> <p>Human health: Increase in malnutrition, particularly among children (deaths of children and elderly were highlighted for 1968).</p> <p>Education: School performance decreases; resource-poor families withdraw children as they cannot cover education-related expenses.</p>
Other	<p>Cholera epidemic 1972: Deaths of children and elderly</p> <p>Earthquake: Destruction of houses</p>	<p>Flood of 2002: Impacts similar to above, but shorter in duration. The community has not been able to fully recover the partially destroyed irrigation channels, which further constrained the planting options during the subsequent drought. Positive effects: Revitalization of the occasional fields and extension of pasture after the flood receded.</p>	<p>Cholera 1992: Deaths of children and adults</p> <p>Earthquakes 1970, 1972: Caused a spilt in the Piura riverbed.</p>	<p>Heavy rains, 2001: Collapse of houses, isolation due to damages to roads and bridges, loss of animals, erosion.</p> <p>Strong winds: Occur annually in May and June, sometimes lasting into July; knock down crops and trees and cause damage to houses and an increase in erosion.</p> <p>Fog: Occurs annually during the rainy season and usually continues until May and June. The persistent humidity promotes the proliferation of pests and hence impacts the income sources of the population.</p>

The challenge for the communities in Piura lies in being able to overcome the destructive effects of El Niño and to utilize its benefits in order to build up assets that carry them through marginally productive dry years. For highland community Cabuyal, the smaller but annually recurring climate hazards represent additional constraints in securing livelihoods. The ability to anticipate and prepare for climate events such as El Niño should therefore theoretically help to reduce the existing vulnerabilities.

During strong El Niño events, the excessive rainfall collecting in the Piura River basin triggers often massive increases in water levels and water volumes transported over time. For example, in 1998 the average level of the Piura river was 721 percent above normal (CAF 2000). During El Niño conditions that developed in December 1997 and March 1998, the river carried the greatest amount of water. The water flow peaked at 4,424 m³/s on March 12, which represents the date when Chato Grande was flooded.

The flooding reported in Chato Grande in 2002 was triggered by two intense rainfall periods in the upper Piura River basin. The first rainfall period occurred April 1–4, and the second April 6–7, while no intense precipitation occurred in the lower river basin at the same time. The water discharge from the highlands into the Piura river resulted in maximum water-flow volumes of 3,750 m³/s (Angulo, personal communication). Consequently, Chato Grande and other communities were flooded when the crops were in full growth.

Concerning the reporting of drought periods, it is important to stress that lower (bajo) Piura is generally characterized by low rainfall. According to the climate profile for the Piura River developed by SENAMHI, the rainfall for the Sechura coast averages 65 mm during a normal rain period (Angulo, personal communication). The water input is therefore minimal considering the high rates of evaporation. It is only during the strong El Niño events that the coastal areas receive significant amounts of rainfall; e.g., 400 mm

from September 1982 to April 1983 and 1,000 mm from September 1997 to April 1998.

Given the general absence of pronounced rainfall during normal years, dry years as classified by the coastal community of Parachique, Chato Grande, and Locuto are predominantly associated with lack of water supply from the Piura River during the rainy season. It is the resulting impact on the agricultural productivity that then leads to the classification of years as drought years. Given that the communities refer to agricultural droughts, this may explain the different start and end dates of some of the drought periods highlighted by the individual communities in the assessments. Also, drought conditions do not always develop uniformly throughout the Piura River basin, as was the case in 1960s, where lower and upper Piura was simultaneously affected. The selectivity of human memory also has to be accounted for, although it can be expected that significant events that directly impact on livelihoods within the communities will be recalled fairly accurately.

The exposure to climatic hazards differs between the lower, middle, and higher parts of the river basin. On the coast, air temperature increases in the months preceding the heavy rains during 1997 further affected flowering and fruit production of crops. In the highlands (sierras), the relative humidity increases. Cabuyal is located in the central sierra of Piura, where the lower height of the Andes allows for moisture transport from the Amazon region. In general, the rainy season lasts for six months, with approximately 80 percent of the rainfall occurring from December to April. During El Niño years, regular moisture and rainfall conditions tend to intensify, but significant rainfall episodes are also possible in normal years, such as 2001 (Angulo, personal communication).

Puno

The communities assessed in the department of Puno in the Peruvian altiplano all express concerns for a

broad array of climatic hazards. Floods, droughts, frost, hail, and snow are all highlighted as concerns in all of the communities (Table 9). The relative importance of the hazard varies by community, which may reflect the different proximities to Lake Titicaca, as well as other topographic characteristics that influence the microclimate. The data is drawn from the community assessment report provided by Valdivia and Quiroz (2006) for this study.

In each community, the participants were asked to indicate the degree of impact associated with each hazard. Impacts could be classified into severe, medium, or little/none. While these are subjective rankings, they give a crude insight into whether these events tend to affect entire communities or individual households. A severe impact is generally associated with entire loss of the harvest, severe impacts on assets, or impact on human health. It represents a disaster on the household level. A medium impact indicates a considerable impact on the asset base, which translates into hardship for the household.

Drought is generally considered a key vulnerability by the communities. Being ranked once as the worst hazard and four times the second-worst hazard the communities are exposed to (Table 9), the impacts tend to be widespread within the community, affecting on average 74 percent (n=265) of the households severely and having a medium degree of impact on another 20 percent of the participants voting in the

assessment. Most severely affected is the community of Santa María. In contrast to this general picture, the community of Alto Aychuyo ranked droughts lower, and community members were more concerned with events associated with cold temperatures, such as hail, snow, and frost. This likely reflects the higher elevation of the community and its greater distance from Lake Titicaca.

Frost is the other hazard ranked highly by several communities. According to the votes of the participants, the impacts are also widespread among the households, affecting 74 percent of the participants severely and another 21 percent moderately. Yanamocco, Cari Cari, and Ancacca considered frosts as their key vulnerability, ranking it higher than all the other hazards.

Floods and hail display diverse rankings when compared across communities and snow events, with the exception of Alto Aychuyo, usually ranked lowest among the five climatic hazards identified in all the communities (Table 9). The impact of hail and floods is less uniform across the communities, affecting on average around 61 percent of the participants (Valdivia and Quiroz 2006). Both hazards, however, show a broad range of the vulnerability from community to community.

While in Aychuyo 94 percent of the participants indicated that they are severely affected by hail, only

Table 9. Qualitative Ranking of Natural Hazards by Communities in Puno

Ranking	Community					
	Candile	Yanamocco	Cari Cari	Alto Aychuyo	Santa María	Ancacca
1	Floods	Frost	Frost	Hail	Drought	Frost
2	Drought	Drought	Drought	Snow	Floods	Drought
3	Frost	Flood	Hail	Frost	Hail	Floods
4	Hail	Hail	Flood	Drought	Frost	Hail
5	Snow	Snow	Snow	Flood	Snow	Snow
6				Wind		Wind

Source: Participatory workshops with community members. CIRNMA-CIP, July 2005.

47 percent of participants in the assessment of Cari Cari indicated that they are severely impacted. Candile, which is at a lower elevation and in the vicinity of Lake Titicaca, is most concerned about floods (with 86 percent being severely affected); hail events are less of a nuisance (62 percent severely affected) than in communities located at higher elevations, such as Alto Aychuyo. Alto Aychuyo and Ancacca also highlighted as a concern their exposure to strong winds.

Spatial Disaster Risk


All the communities highlighted their exposure to multiple climatic hazards. In order to further understand to what extent the spatial exposure to these hazards overlaps, CIRNMA asked communities to describe the approximate area that was severely affected by the individual hazards. Focus groups focused on specific hazard events and described the spatial extent of the impact in relationship to major landmarks. A Global Positioning System (GPS) was then used to develop spatial risk maps for each community. Figure 8 shows an example of a spatial risk map for the community of Santa Maria. The maps for the individual communities can be viewed in detail in Appendix A.

Santa Maria and Yanamocco (Appendix) show that basically the entire region of the communities is exposed to hail events. This is also the case for drought, frost, and snow. Floods are more spatially confined. In Santa Maria, floods result from river overflow, while in Yanamocco it is the combination of river floods and a high water

table. While displaying similar characteristics in terms of overlapping spatial exposure to climatic hazards, the impacts on livelihoods are different. Santa Maria places a greater emphasis on crops, while Yanamocco concentrates on dairy cattle production.

Cari Cari and Ancacca place different emphases on livestock and crops, which may be related in part to different spatial exposure to hazard risk. In comparison, Ancacca has more areas of overlapping events. In addition, most of the land is located in the pampas (flatland). The community focuses on livestock production, as agricultural activities are very risky. While Cari Cari also has pampas, the areas with overlapping hazard risks are smaller. The community has a greater proportion of land under cultivation, while also possessing pastures to support livestock.

Figure 8. Spatial Exposure to Hydrometeorological Hazards of the Community of Santa Maria




While being exposed to the same hazard types, the spatial exposure differs considerably between the communities of Aychuyo and Candile. Aychuyo, which is located near the border with Bolivia at a higher elevation and at a greater distance from Lake Titicaca, is highly exposed to hail. Flood areas are confined to the vicinity of the river. By contrast, large areas of Candile

are exposed to floods due to its flat topography and proximity to Lake Titicaca.

Temporal Exposure to Climatic Risks


In each community assessment in the altiplano, time lines of major climate-related disasters were

Figure 9. Time Line of Major Natural Hazard Events Recalled by the Communities — A



Note: Time line of major natural hazard events recalled by the communities through 1980, including droughts (D), floods (F), frosts (Fr), hail (H), snow (S), and wind (W). El Niño years, as classified by SENAMHI, are represented by the orange bars. Communities may experience a range of hazards in a given year. A lack of synchronicity in the occurrence of hazards between the communities is apparent, suggesting that disasters are fairly localized. Source: Community assessments in Puno.

Figure 9. Time Line of Major Natural Hazard Events Recalled by the Communities — B
(continued)



Note: Time line of major natural hazard events recalled by the communities from 1981–2006, including droughts (D), floods (F), frosts (Fr), hail (H), snow (S), and wind (W). El Niño years, as classified by SENAMHI, are represented by the orange bars. Communities may experience a range of hazards in a given year. A lack of synchronicity in the occurrence of hazards between the communities is apparent, suggesting that disasters are fairly localized. Source: Community assessments in Puno.

reconstructed (Figure 9). While only featuring events that severely impacted the community, these time lines illustrate that the households often face several climatic extremes within a year or consecutive years.

The recollections by the communities in Puno show an absence of synchronicity between most types of disaster events. While drought impacts are more widespread and uniform, the impacts of the other hazards are more localized. This applies in particular to hail events. As a result, it is difficult to ascertain whether the hazard really occurred at this point in time or not, as the disaster reports tend to provide spatially aggregated impacts.

Looking at the OFDA/CRED Emergency Disasters Database (EM-DAT), for example, various droughts are listed as disasters in Puno. This includes the drought of 1982–83, which was listed by five communities. It was associated with a strong El Niño at the time, which caused drier conditions than usual in the entire altiplano region. Interestingly, the strong El Niño of 1997–98 did not have the same effect. The database also lists the drought of 1966, which is a year apart from drought conditions reported from Anccaca. With the absence of local climate data, however, it is not possible to say whether the recollection of the community is off by one year, the reporting is inaccurate, or these are two different events taking place in Puno. It should be kept in mind that communities' responses refer to agricultural droughts, which are visible in yield reductions rather than meteorological drought conditions.

EM-DAT also includes reports of floods in 1984 and 1994 that closely coincide with flood-related disasters reported by the communities of Santa Maria (1984), Candile (1986), Aychuyo (1986), and Cari Cari (1986, 1995). The flood of 1986 also matches the record-high stand of Lake Titicaca (Seimon, personal communication).

Cold spells that occurred since 1990 are documented in EM-DAT for 1991, 2003, and 2004. Only 2004 is featured in the recollection of the communities of Candile and Anccaca. It is interesting that the recollection of frost-related disasters is more noteworthy for events that occurred in the 1960s and 1970s than in recent periods. This may reflect a decreased vulnerability or a lower frequency of events in recent years. In the ranking of hazards, however, frost events are featured as major concerns to the communities. The impact of cold temperatures on children's health and inadequate housing also contradicts this impression.

Overall, the recollection of disaster events show that all hazard types have translated into disasters at the community level over time, illustrating a general adaptation deficit to the prevalent climatic conditions. This reflects the general picture of a high range of climate-related disasters reported for Puno (Table 2).

Impacts

After the identification and ranking of the hydrometeorological risks, the altiplano communities, in focus groups, identified the type of impacts associated with flood, drought, hail, and frost events in the past (Tables 10 and 11).

The recollection of the communities shows a pervasive impact on conditions essential to reducing poverty across disaster types. These include immediate impacts on human health and household assets, as well as impacts on the long-term prospects of advancement of households through loss of schooling time.

As illustrated by the broad scope of impacts, the communities are currently equipped to cope with and adapt to the range of climate conditions they are exposed to. Housing structures are, for example, inadequate for the risk profile of the region, not providing enough insulation against cold conditions and being impacted by flood and hail.

The limited adaptive capacity also becomes apparent in the frequently highlighted livestock sales during times of disasters. Community members resort to short-term

coping strategies that alleviate the immediate impact of the disaster event but deplete their asset base, and hence make them more vulnerable in the long run.

Table 10. Livelihood Impacts of Drought and Floods, Communities of Puno

Sector	Impacts	
	Drought (community)	Flood (community)
Health	<ul style="list-style-type: none"> - Malnutrition (all) - Diseases: cough, fever, stomach, and skin irritations (all) - Increased child mortality (all) - Measles epidemic (SM, Acc) 	<ul style="list-style-type: none"> - Diseases, mainly stomachal and respiratory (all) - Malnutrition, especially children and elderly (Cd, Ym, CC) - Increased child mortality (Ym) - Skin problems and bone aches (Cd, Ay, Acc)
Education	<ul style="list-style-type: none"> - Somnolence, lack of concentration (Cd, CC) - School desertion for work & lack of resources (all) 	<ul style="list-style-type: none"> - School desertion (all) - School closure (Ay, SM, Acc) - Low educational achievement (Cd, CC) - Increase in the time required to reach school (Cd, Ym)
Housing & Infrastructure	<ul style="list-style-type: none"> - House ceiling dries up (all) - Shortage of totora for roof rebuilding (all) - Drying of wells (Cd) 	<ul style="list-style-type: none"> - Destruction of houses (all) - Road destruction (all, except SM) - Destruction of wells (all, except Ay and SM) - Migration (Cd, CC) - Destruction of bridge (CC, Ay, SM) - Latrine destruction (Cd, Ym) - Destruction of community site (Cd) - Destruction of church (Cd)
Agriculture & Livestock	<ul style="list-style-type: none"> - Loss of all crops (all) - Pest outbreaks (all) - Migration of young people and adults looking for work (all) - Food shortage (all) - Shortage of fodder (all) - Increased livestock mortality (all) - Weight loss in livestock (all) - Increase in skin parasites (all) - Price reduction for livestock sales (all) 	<ul style="list-style-type: none"> - Loss of crops (all; with exception of Ay, the communities indicate that the entire harvests were lost) - Increase of weeds - Lack of animal fodder (all) - Increased livestock mortality (all) - Weight loss (all) - Increase in livestock diseases, especially pneumonia and diarrhea (all) - Hoof softening (all, except Ym) - Low livestock prices, difficult to sell (Ym, CC)
Handicrafts	<ul style="list-style-type: none"> - Shortage of wool (Cd, CC, Ay, Acc) - Lack of resources to purchase wool (Cd) - Increased preparation of handicrafts (Ym, SM) - Low price for handicrafts (Ym) - Handworks were not carried out (Ay, Acc) 	<ul style="list-style-type: none"> - Loss of handicrafts (Cd) - Loss/shortage of raw material (all, except Ym) - Reduction of wool quality

Note: Communities: Acc = Anccaca, Ay = (Alto) Aychuyo, CC = Cari Cari, Cd = Candile, Ym = Yanamocco, SM = Santa Maria.
Source: Authors' data, summary of community assessments.

Table 11. Livelihood Impacts of Hail and Frost, Communities of Puno

Sector	Impacts	
	Hail (community)	Frost (community)
Health	<ul style="list-style-type: none"> - Disease: Respiratory, colds (all) - Increased child and elderly mortality (Ym, Ay) 	<ul style="list-style-type: none"> - Disease: Respiratory, colds (all) - Increased mortality of children (all) and elderly (Cd)
Education	<ul style="list-style-type: none"> - Absence from school classes (all) - Repetition of school year (all, except: Ym, CC) - School desertion due to migration (Ay, SM) 	<ul style="list-style-type: none"> - Absence from school or delay of classes (all) - School desertion & migration (all except Cd)
Housing & Infrastructure	<ul style="list-style-type: none"> - Damage of house ceiling (all) - Damage to school (Cd) - Damage to wells (all) 	<ul style="list-style-type: none"> - Low temperatures inside the houses (all) - Damage to the house ceiling (Ym, CC) - Frozen wells (all) - Very low temperatures in school buildings (all)
Agriculture & Livestock	<ul style="list-style-type: none"> - Damage to crops (all) - Increase in plant diseases and pests - Shortage of food - Shortage of seed - Increased livestock mortality (all) - Weight loss of livestock - Increase in livestock diseases and ailments (all) - Low milk production (Ay) - Forced sale of livestock (all) - Livestock theft 	<ul style="list-style-type: none"> - Loss of all the crops (all) - Shortage of food (all) - Shortage of seeds for the next planting cycle (all) - Migration (all) - Increase in pests (CC, Acc) - Freezing of grass, shortage of pasture (all) - Increase in livestock mortality (all) - Weight loss of livestock (all) - Increase in parasites and animal diseases (Ay, SM, Acc) - Low milk production (all) - Forced livestock sale at low price (all)
Handicrafts	<ul style="list-style-type: none"> - Low wool production (all except Ym, SM) - Greater production of woven fabrics (Ym) 	<ul style="list-style-type: none"> - Poor quality of wool (CC, Ay) - Shortage of wool (Acc, Ay) - Greater production of wool fabrics (Ym, Ay, SM) - Smaller production (CC, Acc) - No handicraft production (Cd)

Note: Communities: Acc = Ancacaca, Ay = (Alto) Aychuyo, CC = Cari Cari, Cd = Candile, Ym = Yanamocco, SM = Santa Maria.
 Source: Authors' data, summary of community assessments.

Comparison of Risk Exposure of Communities in Piura and Puno

Comparing the responses from the community assessments in Piura and Puno, it is apparent that the altiplano communities face the challenge of building adaptive capacity to a broad spectrum of climatic hazards with very different characteristics. In Puno, communities have dealt with high daily fluctuations in daily temperatures and exposure to extreme events, such as hail, cold spells, droughts, and flooding. Often, these climatic extremes overlap in space and time.

By contrast, the communities assessed in Piura face climatic extremes that are associated with either too wet or too dry conditions, representing a bimodal risk exposure. Furthermore, the risk exposure is more clearly linked to El Niño and La Niña conditions than in the altiplano, making the challenges more predicable in

northwestern Peru. Another difference is that climatic changes associated with El Niño may not be entirely negative. Depending on the strength of the event, the increase in precipitation and the associated positive effects on ecosystem productivity can offset negative impacts and actually help households to develop their asset base.

The broad range of impacts highlighted by communities in Piura and Puno showcase a general awareness within the communities of their exposure and vulnerability, as well as a limited capacity to do something about it.

Vulnerability

Within the context of this report, vulnerability is understood as the characteristics of the communities

that influence their ability to anticipate, cope, and adapt to climate-related hazards. Both socioeconomic and environmental factors influence conditions of vulnerability. Vulnerability is therefore a dynamic concept, driven by various causal agents and processes (Vogel and O'Brien 2004). Changes in the local environment, market conditions, or climate change can increase or decrease the degree of vulnerability by affecting the exposure to a given hazard as well as the ability to cope or adapt to it. The sections below describe the feedback from the communities concerning their perceptions of which activities, livelihood aspects, and environmental factors affect their vulnerability to the climatic hazards experienced in Piura and Puno.

Socioeconomic Vulnerability

The communities deemed those groups most vulnerable who had the least assets to absorb climatic shocks and limited capacity to spread risks and adjust their income sources to changing environmental conditions.

In general, the communities assessed in Piura and Puno considered livelihoods based solely on agricultural practices most at risk. The communities differentiate groups that are less or more vulnerable according to the amount and quality of land and livestock they own. The area and location of land allows households to decrease the likelihood of catastrophic losses, while the type and number of livestock represents an important additional income source as well as an asset, which allows households to cope better with a climatic extreme.

For example, in the community of Chato Grande in Piura, the most vulnerable group is families of young farmers who possess little or no land and predominantly depend on work as farm laborers. According to the participants in the assessment, this holds true for approximately 20 percent of the population. Also highly vulnerable are large families (eight to nine children) that own only small patches of land (1/2 ha or even less). This group is also dependent on complementary income as farm laborers. If their small

area of land is lost, there is very little opportunity for this group to recover from the shock, as farm laborers tend to lose their jobs in the times of crisis.

In addition to the size of the land, the community of Cabuyal in the sierras of Piura and the communities of the Peruvian altiplano noted that the vulnerability is influenced by the ability to plant at different altitudes.

The possession of livestock is another factor that influences the vulnerability within the community. In general, the communities possess only limited monetary assets, livestock constitutes their “checking and savings account” (R. Quiroz, personal communication; Valdivia 2001). It also represents an insurance against climatic shocks, enabling households to ensure food security, acquire seeds, and building material. In the assessment, groups that possess only small numbers of livestock were considered more vulnerable.

In Piura, resource-poor families raise domestic fowl, pigs, sheep, and goats, but do not possess cattle. It was pointed out that households owning larger numbers of livestock may lose a greater number of animals in a flood, but poorer households are in danger of losing their entire livestock. Thereby their asset base for paying for the education of their children and recover from future shocks is depleted medium- to long-term, reinforcing conditions of poverty.

Another aspect of vulnerability raised in the community assessments was the ability of household to adjust their livelihoods to the prevalent environmental conditions. For example, in Parachique community representatives considered households that only live from the dry forest most vulnerable. For these households, food security and income depends on the ability to grow crops for flour and cereals from sparse rainfall during the rainy season.

Families in Parachique, which use the marine resources during the normal and La Niña years and incorporate farming activities in their activities during El Niño years, are considered generally less vulnerable. Among

the fishermen, however, different levels of vulnerability were identified. The most vulnerable groups are the Pinteros, who possess only small boats and fish on a small scale using hooks and lines. Their catch is sold for local consumption. In Parachique, Pinteros are either young men who are just beginning to learn to fish or immigrants.

For communities in Piura, which are located inland, the ability to combine income activities from agricultural and fishing activities is not an option. Therefore, the communities define vulnerability largely according to land access, size, and quality and the possession of livestock. Alternative income sources are comprised of labor in nearby cities. Income can also be replenished during rainy years by expanding agricultural productivity to the occasional fields.

In the altiplano, the communities highlighted handicrafts and informal trade as alternative income sources. Tables 10 and 11, however, show that in some communities these alternative income sources are impacted adversely at the same time as the main livelihood activities and hence play only a limited role in reducing the impact of climatic shocks on the assets of a household.

Environmental Vulnerability

In Piura and the altiplano, the compounding effect of environmental degradation and change with the vulnerability to climatic shocks was recognized.

In Parachique, community members highlighted inadequate health and hygiene conditions and the pollution of the ocean as major problems, as it has reduced fish stocks and also limited their ability to sell fishing products on the market. These environmental conditions therefore compound the adverse impacts of El Niño events on the fishing sector and constrain the ability of the community to acquire monetary assets. In alto Piura, the community of Cabuyal recognized the conflict between land demand for agricultural activities

and the increase of erosion due to clearance of forest cover. While the community recognizes deforestation as a short-term solution, it also feels there are few alternatives when fertile soil is lost due to intense precipitation or strong winds.

In the altiplano, overgrazing of land resources is a recognized problem, but during the workshop in Lima, community members expressed particular concern over changing climatic conditions and dwindling water supplies, which further increases the uncertainty and vulnerability of their agricultural practices. This will be discussed in further detail in the section on perceived changes in climatic conditions.

Long-term Consequences of Vulnerability

A shared concern of the communities in Piura and Puno is the vulnerability of the children and youth to climatic risks. In the altiplano, malnutrition and increased child mortality are highlighted by community members for multiple climatic hazards. Aside from the concern for the health of the children, it is also the long-term consequences resulting from the climatic hazards that are emphasized. Reduced schooling time has been cited both on Piura and the altiplano. There is the strong perception that this limits the opportunities of the children for the future. For example, in the community assessments it was stressed that with their education, community children cannot compete with children from larger cities, and therefore the existing, risk-exposed livelihoods represent their only option.

Key Aspects of Vulnerability

The community assessments reveal a keen awareness among the participants about which factors contribute to the vulnerability to environmental risks (Table 12). There is also the recognition, however, that some of these conditions of vulnerability are beyond their control.

Table 12. Conditions of Vulnerability Highlighted by Communities

<i>Piura</i>	<i>Puno</i>
Livestock: number and quality	Livestock: number and quality
Land: ownership, access, size, quality, location	Land: ownership, access, size, quality, and location; i.e., ability to plant at different altitudes
Access to alternative income sources: fishing (coast), handicrafts, trade	Access to alternative incomes sources: handicrafts, informal trade, fishing
Environmental degradation: pollution (coast), deforestation, erosion	Environmental degradation: overgrazing, erosion Environmental change: Climate, Water

Source: Authors' data.

While land ownership, and the quality and size of land are recognized influences on a household's resiliency to climatic shocks, land-poor households have very little options to change this condition. Even in Chato Grande, the access to irrigation is only of limited benefit to land-poor households. Water rationing constrains the productivity of the already limited land resources, while at the same time the demand for farm labor also decreases. Hence, the self-sustaining capacity of the households is further constrained.

Given that dry conditions in Piura are largely linked to La Niña conditions, which often precede or follow wet El Niño years, these households are exposed to prolonged periods of stress, which make it difficult to maintain or build assets necessary for education or investments.

In the altiplano, the frequent exposure to a variety of climatic extremes means that households that own little land and livestock are in constant danger of losing their entire basis for livelihoods. Once all livestock is lost, it is difficult to acquire new animals without access to credit, leaving the household more vulnerable to further climatic risks.

The interaction between environmental degradation and vulnerability to climatic hazards also illustrates the limited ability households have in reducing their long-term vulnerability. In bajo Piura, as illustrated by the community assessment of Cabuyal, intense precipitation leads to the erosion of topsoil. This,

in addition of the demand for fuelwood, promotes deforestation activities, which then in turn further exacerbates erosion processes and the capacity of the land to retain water. In addition, the downstream transport of sediment raises the riverbed and increases the flood risk in lower Piura.

Conditions of vulnerability and resulting choices in land-use and resource-use management can impact other locations as well. As highlighted above, the excessive use of natural resources in the Noma microbasin is compounding water-shortage and flooding problems downstream. While in Piura (Catacos and Sechura), the construction of dams has further increased the silting of riverbeds, they often collapse during intense rainy seasons and hence do not reduce the flooding risk (Angulo, personal communication). Therefore, the vulnerability of one location not only influences the choices of households by favoring short-term over long-term solutions, these choices can also affect the risk exposure of other communities, which are connected through their mutual location along a river's course.

Coping and Adaptation Strategies

Socioeconomic conditions of vulnerability as well as driver effects of environmental degradation are the outcomes of the coping and adaptation capacity of households and communities. The range of options available to anticipate risks, take risk-mitigating measures, and reduce the impact of a given extreme

event will influence the asset base and opportunities for advancement of households (Valdivia et al. 2003).

In addressing climate risks, one can distinguish between coping measures, which are focused at mediating the specific impact of a hazard that has occurred, and adaptation strategies, which are focused on systematically reducing underlying vulnerabilities to hazards with the goal of avoiding disastrous impacts.

Coping Strategies

While the disaster risk profile in Northwestern Peru and the altiplano differ in many ways, coping mechanisms highlighted in both areas during the community assessments show considerable commonalities. In both regions selling livestock and other assets and migration represent pervasive coping mechanisms to dampen the magnitude of a disaster event at the household level.

Selling of Livestock. In the face of climatic shocks, a common coping strategy is to sell livestock to obtain essential food supplies and recover seeds for the next planting season and construction material. The possession of livestock represents, therefore, an insurance to households, when harvests fail and investments are required for recovery. As a coping strategy, the selling of livestock is associated with several caveats.

The assets of farmers in the communities largely consist of the livestock they own. Monetary income tends to be spent immediately on food and education. If there is a surplus, this will be used to improve housing conditions, obtain livestock, or diversify activities.

The sale of livestock in response to a hazard event is a strategy shared throughout the communities. The communities of the altiplano note that the various climatic hazards are all associated with a drop in the sales price. Because many community representatives employ this coping strategy at the same time, there is a high supply of animals, which in times of climate

shocks are often also malnourished, resulting in low prices. It therefore reduces the immediate impact of the climatic shock but also depletes the asset base of the household.

Where families are forced to sell their entire livestock, this solely represents a short-term remedy, which likely increases vulnerability in the medium to long term to subsequent climate risks, as the household becomes more dependent on more climate-sensitive agricultural activities (if it cannot replenish its assets by acquiring livestock or diversifying income sources).

Migration. When harvests are destroyed and the household has no ability to liquefy assets to recover, migration of household members often represents the only option to support the family. Migration is not really a choice, but an option of last resort. In Piura and Puno, the communities highlighted the disruption of the social networks and family structures as well as the impact on the educational opportunities of the children.

When households lack alternatives, household members or entire families migrate in search for jobs to industrial centers at the coast or the Amazonian lowlands. Table 13 summarizes the migration destinations of members from the communities assessed in Piura and Puno, showing that the destination also sometimes depends on the type of hazard. Droughts and floods have different impact durations and hence pose different requirements for alternative income sources.

The lack of education and marketable skills forces the farmers to take low-wage jobs in unfavorable working conditions, but this remains the only option to compensate for losses and ensure the food security.

Migration is unwanted and does not really represent a choice, but rather a lack of alternative income opportunities. Communities highlighted the adverse effects on family structure, social networks, and local knowledge.

Table 13. Destinations of Push Migration, Piura Communities

Community	Main Destinations
Parachique	Flood: Following the 1982–83 floods, part of the community relocated to Ciudad del Pescador, a settlement nearby
Chato Grande	Drought: Tumbes, Sullana, Chepen, Huaraz, Lima Flood (more temporary): Lambayeque, Chiclayo, Chimbote, Lima
Locuto	Drought (men in search of farm labor): Bajo (lower) Piura, Chiclayo; (women working as maids): Tambogrande, Piura (city)
Cabuyal	Drought: Towns in the Peruvian Amazon, e.g., Jaen, Nuevo Cajamarca, San Ignacio, Saramerisa, Pueblo Nuevo de Maray

Source: Authors' data.

Adaptation Practices

Adaptive measures taken by humans to address climate risks can be anticipatory and reactive. In general, adaptation leads to a reduction of vulnerabilities. Ideally, adaptive measures should prevent climate hazards from turning into disasters and enable stakeholders to maintain and develop their asset base. In contrast to coping measures, adaptation in the context of this paper should be considered as a planned response that is informed by early warning of extreme events, traditional knowledge and practices, and/or intrinsic understanding and evaluation risk factors and remedial options, which are also influenced by available assets. In contrast to coping measures that mitigate the near-term impact of a given hazard event, adaptation measures should ideally lead to sustained vulnerability reduction of the household and community.

Diversification of Income Activities. This is crucial in reducing vulnerabilities to climatic risks. When discussing aspects of vulnerability, it was, for example, noted that coastal communities such as Parachique households that combine fishing and agricultural activities tend to be less vulnerable than the ones who only focus on agriculture (impacted by droughts and floods, benefiting from extra moisture input during El Niño) or artisan fisheries (impacted by El Niño).

In the altiplano, communities supplement their income through handicrafts and weaving. When highlighting impacts, however, in the majority of the communities these activities were often also affected

by the disaster event due to a drop in sales price or limited supply of raw material, such as wool. Only two communities, Yanamocco and Santa Maria, did not indicate an impact on handicrafts dependent on wool supply. Yanamocco residents highlighted increased production during these events. This may suggest that in Yanamocco (and Santa Maria) there is some foresight planning, ensuring sufficient supply of raw material. By contrast, the other communities face dwindling supplies in the aftermath of the events when there would be a need to compensate for losses in livestock and agricultural productivity.

Livestock Management. Livestock has been already been highlighted as an important coping strategy. The possession of livestock becomes an adaptation strategy when it allows the household to minimize the impact of climate risks to maintain and expand its asset base. The household may sell some of its livestock in times of disaster, but possess enough animals to subsequently increase its livestock size.

In Piura, community members pointed out that less-vulnerable households succeed in maintaining a core livestock throughout flooding events in El Niño years and are then able to increase livestock numbers when the positive effects of increased pasture size materialize in the wake of the additional moisture input. As a consequence, El Niño turns into a benefit, allowing these households to increase their livestock numbers and hence their assets for investments in housing, education, or health care.

Households that can maintain livestock throughout extreme climatic events are able to sell animals in an anticyclical manner, at times when there is greater demand and hence better prices. Livestock products such as milk and cheese further supplement diet and income during this time.

The challenge for households lies in maintaining a critical number of livestock throughout the occurrence of climatic hazards. In Piura, this means ensuring livestock health through flood and drought conditions, while in Puno livestock needs to be protected against floods, drought, frosts, and hail and associated implications for food security. In the altiplano, the number of livestock per household roughly increased with altitude and distance to water bodies such as Lake Titicaca, which buffers against climatic fluctuations and offers some protection from cold spells for agricultural productivity.

In the altiplano, the communities also recognize the importance of protecting the livestock against not only extreme events, but also high diurnal temperature variations. As will be discussed in the recommendations, research has shown that gains of livestock productivity and, hence, the ability to increase income, are substantial when using shelters and in combination with improved animal feed. Despite this recognition, feedback from community members suggests that investments in the construction of shelters is too costly without access to affordable credit.

Expansion of Agricultural Activities. In Piura, communities utilize the added ecosystem productivity during El Niño years, expanding planting areas and exploiting products of the revitalized dry forest. In the assessments, the different options available to households depending on their affluence (access to seeds, fertilizer, land, number of livestock) was highlighted.

According to responses from community representatives, resource-poor households will focus

on making an immediate profit from the improved productivity. More affluent households can afford to wait, processing agricultural products and then selling them at higher returns later.

In the altiplano (as well as Cabuyal), farmers focus on spreading risks by planting at different altitudes and using a variety of crops. This option is, however, only available to the more affluent groups in the communities.

Traditional Adaptation Measures. The assessment in Piura did not reveal the use of traditional water-harvesting structures or cropping systems to address existing environmental risks. This stands in contrast to reports from coastal communities in Ecuador (Marcos, personal communication), which face a similar risk exposure (see appendix F for further information on Ecuador). In coastal Ecuador, communities still use *albarradas*, ancient structures that collect excess water during El Niño years. Being built as u-shaped dams on semipermeable soil, the water percolates, recharging the aquifer. This allows for better water supply during dry conditions, turning aspects of El Niño into an advantage.

The absence of such structures in the communities in Piura may in part be explained by the history of the region. Large-scale farming predominated until, during a land revolution, the land was divided and distributed to laborers. The previous large-scale production for export may not have favored the existence of these structures and hence led to their destruction. Another possibility is that the drier climatic conditions did not favor these water-harvesting structures as much as in coastal Ecuador. It may be worthwhile to explore whether the traditional water-harvesting techniques could also be of relevance to managing agricultural production risk in Piura.

In the communities of Candile and Yanamocco, “raised fields,” also called *waru warus*, are still in use. *Waru warus* consist of mounds of fertile soil, which

are intersected by small water channels. The resulting structure creates a microclimate that provides moisture and also buffers against temperature fluctuations. Raised fields are also found in areas of northwestern Peru (although they have not been documented in the community assessments in Piura; Quiroz, personal communication), but more elaborate and permanent structures are characteristic of the altiplano highlands. Other structures aimed at adapting to prevailing climatic conditions include sunken fields (coastal dry areas) and conchas (highlands).

In the altiplano, traditional social networks also play an important role in mitigating the impact of climatic risks on households. The *aynoka* system represents a traditional decision-making process, where community members decide what crops are planted and where, but farming activities are then carried out individually. This allows for a collective assessment of risk and additional guidance for planting decisions. It may also provide a valuable entry point for external climate information. In the community assessments, Santa Maria and Alto Aychuyo use the collective decision-making process associated with the *aynoka* system.

In times of crisis, *Anyi* is used as a traditional communal mechanism, where access to food and housing is provided in exchange for communal work. Community representatives in Yanamocco, for example, highlighted *Anyi* as a response mechanism to climatic impacts.

External Assistance and Barriers

The external support highlighted by the communities in Piura was predominantly focused on coping with particular disasters. Exceptions are the communities of Parachique and Chato Grande, where the hazard-risk exposure and disaster events triggered efforts to relocate the communities.

The bay area, where Parachique lies, was classified as a high-risk area by Civil Defense due to its vulnerability

to El Niño-related flooding, storm surges, and earthquakes. Following the destructive impacts of the 1982–83 El Niño, the settlement Ciudad del Pescador was created on an elevation nearby and the population was urged to permanently relocate.

The majority of the population, however, returned to Parachique. Reasons highlighted in discussions with community representatives included the fear that boats and fishing equipment would be stolen. While Parachique residents own homes in Ciudad del Pescador, the majority does not live there permanently now.

The risk classification has caused resentment toward authorities, because it limits the use of public funds for improving the infrastructure in the community and hygiene standards in the bay. The resettlement promoted by the authorities is not recognized by the community as an initiative for them, but as an action against them.

In Chato Grande, houses were frequently destroyed during flooding events associated with the strong El Niño events during 1982–83 and 1997–98. With the encouragement of families relocated to a new settlement called Nuevo Chato Grande, institutions provided reservoirs for water supply and latrines in each house (ITDG 1998). It was highlighted, however, that many community members use dual residencies as a strategy for adapting to the prevalent climatic conditions, rather than relocating permanently to the safer location of Nuevo Chato Grande. Others have chosen to remain in Chato Grande, despite being aware that this old settlement is more risk-prone. Obstacles cited by the community members to permanent relocation are the absence of electricity and development projects, as well as the greater distance to cropping and herding places. As in Parachique, living away from the livestock and workplaces increases the danger of losing assets through theft.

In the altiplano, the communities highlighted a range of development activities that would also strengthen their

resilience to climatic shocks. This includes, for example, technical assistance in livestock management, animal husbandry, and handicrafts by the Corredor Económico Puno-Cusco for the communities of Ancacca and Yanamocco. Caritas is supporting Yanamocco in setting up cultivated pastures and, with the Ministry of Agriculture, provides training in animal production and health. Caritas also works with the community of Accopata in establishing animal forage (now in collaboration with the Municipality of Mañazo). CIRNMA works on credit and microfinance schemes with the communities of Accopata and Yanamocco, and also provides support in livestock management. Other organizations that have worked with the communities include PRONAMACHS (Alto Aychuyo, Santa María), PECSA (Accopata), INTERVIDA (Alto Achuyo) and PRADERA (Alto Achuyo).

The responses from the communities showcase the great variety of external support focused on improving livestock health and productivity and providing frameworks for communities to strengthen their asset base. The responses, however, also illustrate differences in the access of communities to development organizations. The activities are usually very specific in their scope and not part of integrative framework; rather, they reflect the specialization of the organizations.

When the community representatives were brought together to discuss the findings of the individual community assessments, it was the first time they had met. The information exchange and hence the learning benefit from successful project activities is thus very limited.

Concerning the support of government institutions, community members expressed their confusion about who to go to in times of crisis. The governance structure and support mechanisms appeared not to be sufficiently clear.

A shared concern emerging in the discussions with communities in Piura and Puno is the lack of access to credit. In Parachique, the fishermen had access to a development fund called Fondo Nacional de Desarrollo Pesquero (National Fund for Fisheries Development), which helped them to finance improvements to boats and fishing equipment with low-interest loans. This fund is no longer available, leaving especially the most vulnerable group, the artisanal fishermen, without viable financing options. In the altiplano, community members complained about the access to agricultural banks. In their absence, the microfinance initiatives piloted by organizations like CIRNMA can benefit individual communities by providing important insights into lending to households in high-risk environments, but they cannot provide widespread relief to the region.

External Information—Weather Forecasts, Early Warning, and Seasonal Climate Outlooks

Within the context of adaptation, timely release of climate forecasts as well as warnings can help communities minimize damages by allowing them to take risk-mitigating measures. The community assessments explored to what extent individual communities have access to information and whether they trust and respond to available information.

Both communities in Piura and Puno receive climate forecasts through the radio, which is the preferred medium. Low levels of literacy and limited access to TV make other media transmission channels less favorable.

El Niño forecasts are predominantly utilized in northwestern Peru. Whether adaptation measures are undertaken, however, depends largely on the alignment of the forecast with local environmental indicators, such as excessive temperatures and relative humidity in the months preceding El Niño events. If there is a disagreement between the forecasts and local indicators, communities tend to follow the indicators. It is also

important who transmits the forecasts—government institutions are generally not well trusted, while certain radio broadcasters have credibility with the community (despite the fact that they basically use the same source of information, e.g., SENAMHI in Peru).

Another problem is the late release of El Niño forecasts to public channels that reach the communities, a delay largely attributable to political and economic reasons. The public declaration of an El Niño event triggers emergency response programs, which are costly. Therefore, there is a hesitation to release forecasts early.

Access to and Use of Information in Piura

While all of the communities have access to weather and climate information, the trust placed in external information sources is generally low (Table 14). Local sources receive more attention and are usually given priority when responding to warnings.

In all communities, the most important medium for outside information is the radio (see Table 14 for information on stations). TV and print media represent additional sources, but are limited in their reach. The access to written information in all communities is constrained by low literacy. The highland community of Cabuyal also has no regular access to newspapers, which can only be obtained during visits to nearby towns.

In the discussions, it became apparent that the communities do not differentiate between the accuracy of weather and climate forecasts. If the experience with weather forecasts is not positive, this will also affect trust in longer-term climate warnings.

Aside from the content, the messenger of weather and climate information is also important. Trust in government agencies is generally low. For example, official warnings from SENAMHI and Civil Defense are questioned in the community of Parachique. Part of the reason is that Civil Defense classified the bay as a high-risk area, which makes it difficult for the

community to obtain public funds. As authorities are perceived as not being on the side of the community, members of Parachique also question whether the weather and climate warnings are intended for their benefit.

In contrast, great trust is placed in the Parachique Harbormaster's office, which transmits information on ocean conditions and warnings of extreme weather via loudspeakers and radio. Experience has shown that this information is reliable and is furthermore aided by the local presence of the office. In the case of conflicting messages from SENAMHI and the harbormaster's office, the community tends to disregard information from SENAMHI, which in their perception is less reliable and not in tune with the local environmental conditions.

Presenters on radio stations also receive more attention than government agencies, although these are the original source of information. The radio programs present the message in a more accessible language to the communities, in addition to being popular and not perceived as driven by ulterior motives. Abraham Levy of Radio Programas del Peru was cited as an important information source.

Concerning El Niño warnings, the community of Chato Grande displayed frustration with the uncertainty and late timing of the forecasts. Community members highlighted the confirmation of an evolving El Niño in 1997 being postponed several times; they heard conflicting messages and when it was finally confirmed, the impacts were already visible.

In addition, the case of Chato Grande also illustrates the importance of a match between forecasts and local environmental signs. In 2002, the community received flood warnings from the authorities. Most of the community, however, did not believe the warnings, because floods usually only occur when there is heavy rain. This was not the case for the first time in the experience of the communities in 2002, as the Piura

Table 14. Access to Weather and Climate Information

<i>Community</i>	<i>Information Source</i>
Parachique	<p>Parachique Harbormaster Type: Ocean conditions, extreme weather alerts, short-term Trust level: High Broadcasting medium: Announcements through loudspeakers and radio access on boats Comment: Based on past experiences, the communities consider this information reliable. This is further aided by their local presence. They are likely to respond to this more than other outside alerts. As the harbormaster retransmits information received from the Callao Harbormaster's office via the Paita Harbormaster, there is sometimes a delay in the alert, which affects preparation time.</p>
	<p>SENAMHI Type: Weather and climate, including El Niño Trust level: Low Broadcasting medium: Radio, TV Comments: Skeptical about information because it has been wrong in the past.</p>
	<p>Civil Defense Type: Weather and hazard alerts Trust level: very low Broadcasting medium: Radio, TV Comment: Community does not trust Civil Defense, which classified the bay as a high-risk area, resulting in constraints in obtaining public investment. The prevailing sentiment is that the institution is not on the side of the community, which also affects the trust level in the potentially beneficial hazard alerts.</p>
	<p>Radio Programs: Radio Programas del Peru, Radio Cutivalu Type: Weather and climate forecasts, including El Niño Trust level: Low Comment: The medical facility transmits radio alerts through loudspeakers. This information was most noted by the women participating in the assessment. There is the perception that the information content is uncertain and the format is unclear, and there is confusion about the time horizon of the forecast.</p>
Chato Grande	<p>Radio Program: Radio Cutivalu Type: Weather and climate information, including El Niño Trust level: Low Comment: There is the perception that climate information is provided late. The participants pointed out that confirmation of the impending 1997–98 El Niño was consistently postponed, which created confusion. When it was finally confirmed, they were already experiencing its effects. This perception also carries over to weather alerts, where there is the feeling that the information is provided with shorter notice than necessary.</p>
	<p>Irrigation Committees, Civil Defense Type: Direct source of information on future climate conditions and practical implications for irrigation Note: The irrigation organizations control the crops that will be grown according to current and projected water availability. When there are water shortages, areas for rice cultivation will be restricted or prohibited. Growing of corn and other less water-intensive crops is promoted. When floods are likely to occur, the irrigation committee does not authorize planting in the floodplains. In conjunction with Civil Defense, information on climate conditions is provided from January to April. Civil Defense regional authorities also visit the community to warn of imminent floods that require evacuation. Comment: Warnings, such as to not plant in floodplain fields when there is the danger of floods, are often disregarded out of economic necessity or when they do not match local environmental signs (as was the case in the flood of 2002).</p>
	<p>Other: The Chato Grande Defense Committee receives warnings and disaster-prevention material from INDECI, the regional government, but the format is not easily accessible. Farmers with schoolchildren pointed out that information they received from the local school was useful, when teachers conducted flood drills with the children, which helped to create general awareness.</p>
Locuto	<p>Radio Programs: Radio Programmas del Peru, Radio Cutivalu Type: Weather and climate information, including El Niño and hazard alerts Trust level: medium Comment: While the community generally places little faith in the weather forecasts, they noted that they received information of the impending El Niño in July 1997 (with the rains beginning in January 1998). The broadcaster they identified specifically was Abraham Levy. It was also noted that Radio Cutivalu broadcasted a flood warning in the El Niño rainy season in 1998 about 12 hours before it occurred. Hence, the community has experienced the benefits of listening to the forecast.</p>
Cabuyal	<p>Radio Programs: Radio Programmas del Peru, Radio Cutivalu, Radio Ronderala Voz de las Alturas (local broadcast from Chalaco) Type: Weather and climate information Trust level: Low to medium Comment: Abraham Levy is noted as a forecaster of Radio Programmas del Peru. The information is considered easily accessible and helps people to stay alert, but it was noted that only a fraction of the community listens to it. Extreme weather warnings are, however, passed on from neighbor to neighbor when being received through the radio, as was the case when they received warning of El Niño, as well as alerts of heavy precipitation in 2001 through Radio Cutivalu. In general, it is more customary to base planting decisions on observations of environmental indicators, and all climate information is compared with local knowledge.</p>

Source: Authors' data, summary of community assessments.

river flood was solely triggered by heavy precipitation in the highlands.

In the highlands, Cabuyal noted the positive experience with earlier warnings of heavy rains in 2001, which proved to be accurate. In contrast with Chato Grande, the community also received El Niño warnings in June 1997, well before the rains started in January 1998. The community members, however, also noted that climate information generally plays a small role in their planting decision, as they tend to trust their reading of the local environmental conditions in judging the quality of the planting season.

Weather and Climate Information in Puno

Given the complex hazard risks communities face in the altiplano, it was of interest to the assessments to gain insights into the extent communities relate to El Niño events and whether they respond to weather and climate forecasts.

The assessments carried out by CIRNMA suggest that almost half of the total participants have heard

of El Niño events. Yet the majority of the community participants relate alerts of El Niño predominantly to the northern coast and not their own situation in the altiplano.

Concerning climate forecasts for the altiplano region, the communities were presented with forecasting formats from SENAMHI to determine whether they trust the information and consider it useful. The responses show that around 47 percent (Table 15) listen to the forecasts. Of this subgroup, 59 percent trusted the content of the information, and a slightly smaller percentage (54) said they would also act upon this information.

The assessments, however, also displayed a great disparity in the degree of trust of external information. Participants from Santa María and Ancacca displayed they highest proportion of knowledge and trust in climate information and willingness to act upon it. These communities had previously participated in the NOAA Andes Climate Variability Project in 2000 and 2001 (Valdivia et al. 2003), in contrast to the other

Table 15. Awareness, Use, and Trust of El Niño Information in Six Puno Communities, 2005 Workshop

Community	Total	Main Information Source	Did Listen	Did not Listen	USE		Trust		Did Not VOTE
					YES	NO	Yes	No	
Aychuyo	42	Radio	12	30	4	7	4	7	0
Santa María	46	Radio	23	21	16	6	14	9	2
Yanamocco	31	Radio	9	13	5	4	5	4	9
Ancacca	56	Radio, TV	31	14	17	14	20	11	11
Candile	53	Radio, TV	27	12	12	15	13	14	14
Cari Cari	37	Radio	22	3	13	9	17	5	12
TOTAL	265		124	93	67	55	73	50	48

Note: Favorite Radio Stations: Pachamama (Puno, subregional), Onda azul (Puno, subregional), Campesina Juli (subregional), San Gabriel (Bolivia, regional/altiplano), Radio Programas del Peru (Lima, national).

Source: Authors' data, based on community feedback.

communities that had received training in using the existing climate information formats.

For the majority of the participants, it is difficult to relate the information to their decision-making processes. These results reiterate the findings of similar studies conducted in Bolivian Aymara communities (Espejo and Carrillo 2000) and in Peru (Claverias), which identified the need to more closely link external information to local knowledge and information networks.

Reflections on Community Responses

Piura. In the responses of communities in Piura, the alignment of local environmental signs with El Niño forecasts was highlighted as important to accepting the message of the forecast. Furthermore, the communities only recognize strong El Niño events as disasters, which have led to widespread damage associated with flooding. Therefore, the communities perceive the recurrence time of El Niño events to be less frequent than the scientific classification of El Niño events.

As shown earlier, strong El Niño events lead to prolongation of the rainy season, as well as an increase in the amplitude of the total rainfall amounts. Therefore, changes in the timing of the rainfall are generally a good indicator of El Niño events and hence reflect the perception of the communities.


Figure 10 relates SST anomalies in the NINO 3.4 region to local changes in temperature and precipitation averaged over the three-month period from January to March (JFM). Substantial increases in precipitation are associated with different SST anomalies, depending on the lead time given. For example, two out of three JFM precipitation

values were above 100 mm when the temperature SST anomaly exceeded 1.5°C in the same period. The temporal relationship between SST anomalies and seasonal precipitation is further displayed in Appendix C.

The important aspect here to consider is that the thresholds for substantial rainfall amounts are higher than the thresholds used for classifying El Niño events. Hence an El Niño event may be classified by scientists focusing on anomalies of the NINO 3.4 region, but not necessarily translate into substantial differences in the environmental conditions of the communities in a specific locale. Consequently, the anchoring of global-scale predictions in local-scale observations becomes important.


The importance of local changes in SSTs in triggering substantial rainfall in coastal northwestern Peru is shown in Figure 11. Here, temperatures above 26°C are associated with elevated rainfall for the month of January, for example. Temperatures above 27°C show substantial increases; this reflects the practice of coastal communities such as Parachique of using local ocean warming as an indicator of the quality of the rainy season.

Figure 10. A Comparison of the Seasonal Precipitation Total from January to March (JFM) for La Esperanza with JFM SST Anomalies in the NINO 3.4 Region



Note: La Esperanza, lower Piura (04°55'04"S, 81°03'38"W, 12 m a.s.l.). Relationship between three-month precipitation totals and SST temperature anomalies in the NINO 3.4 region. While seasonal temperature anomalies of 0.5°C in the NINO 3.4 region are generally used to classify ENSO events, only anomalies above 1.5°C translate into substantial temperature increases in the coastal region of northwestern Peru. Source: Authors' analysis.

Figure 11. Comparison between January Precipitation in La Esperanza and SST Values of the NINO 1.2 Region



Note: SSTs above 26°C are viewed as indicative of local El Niño conditions in northwestern Peru.
Source: Authors' analysis.

Accordingly, the national meteorological service uses a variety of indicators to forecast El Niño events, which includes the NINO 3.4 and 1.2 regions, as well as measurements of coastal SST and land-surface temperatures. Based on conversations with SENAMHI staff, the NINO 3.4 is used as a preliminary alert of possible impending El Niño conditions. The NINO 1.2 region, as well as other observational data, is then evaluated to confirm whether local El Niño conditions are evolving closer in time to the rainy season.

The problem of late confirmation of El Niño forecasts is recognized in part as an outcome of the due diligence process within the ENFEN system, which requires a consensus view. While a uniform message on ENSO conditions is desirable, there appears to be a trade-off concerning the timing of the forecast during this consensus-building process.

At the same time, the information is complementing international announcements and Web postings on the status of El Niño, which may use different classification systems. For example, international institutes such as the IRI use SST thresholds that result in a 25%-50%-25% distribution of La Niña, normal, and El Niño conditions. These thresholds are, however, not necessarily relevant to changes in climatic conditions in northwestern Peru.

Despite the widespread influence of ENSO on climatic conditions, there is no uniform definition of El Niño conditions that are also automatically apply to local climatic conditions. In the case of Peru, the experience developed by national and regional meteorological organizations in identifying thresholds that are locally relevant is of considerable importance. SENAMHI and other organizations do evaluate the accuracy of forecasts. Much

of this knowledge and information on thresholds, however, is not published internationally or made available to the public.

Puno. In contrast with northwestern Peru, it is particularly the NINO 3.4 region that exerts an influence on the climatic conditions of the altiplano. While El Niño in the altiplano tends to be associated with an increased likelihood of drought conditions, these effects may be modulated by other local and regional climatic factors. Furthermore, the multihazard environment in the altiplano means that a variety of climatic extremes may be experienced in any given year. This masks the signal of El Niño events and may explain why the communities in the altiplano consider El Niño events more an issue for the coastal regions in the North.

While advancement in seasonal forecasting techniques and dissemination methods may also yield benefits to the communities in the highlands, awareness and understanding of the multihazard environment already exists. Hence, a strong emphasis should be placed on better integrating this awareness in the design and implementation of development activities to reduce the documented vulnerabilities in housing and infrastructure, human health, livestock health, and management and agricultural activities.

Use of Local Knowledge in the Prediction of Climatic and Environmental Changes

The assessments also included an inventory of physical and biological indicators used within the communities to warn of impending weather extremes or forecast the planting season (Table 16). Some of these indicators are based on long-term observations of the abiotic and biotic environment, while others are rooted in belief systems and superstitions. A challenge is to separate reliable indicators with good a correlation of climate events from indicators which are counterproductive. This is particularly important in relation to establishing better entry points for official climate information, given that the communities tend to trust more information originating from the immediate environment than from outside sources. The communities also expressed concern with some of their indicators becoming less reliable due to climatic changes and an erosion of knowledge due to migration.

Piura

Reflecting the predominant hazards in Piura, all communities highlighted environmental indicators for forecasting wet or dry conditions. The composition of indicators varied considerably between the communities. The participants from the coastal community of Parachique, where livelihoods are largely focused on fishing activities, only focused on the description of physical indicators. The other communities also highlighted observations of plants and species as important signals for judging climate conditions (Table 16).

Cabuyal, as a community located in the highlands of Piura, differed from the other communities in its more-complex risk profile. This was also reflected in a broader use of indicators for forecasting weather and climate conditions (resembling more the importance of indicators).

The trust levels in local knowledge appeared to vary between the communities. In Parachique and

Cabuyal, participants place considerable trust in their interpretation of environmental clues as warnings of weather extremes or insights into seasonal climatic conditions.

When participants in Chato Grande were asked about their use of environmental indicators, it was referred to as a practice of the elder people, viewed as something of the past. In contrast to the other communities, Chato Grande has access to irrigation, making the community less vulnerable to climatic fluctuations. Hence, the reasons for this somewhat dismissive response toward traditional indicators may reflect the perception that such knowledge, or admitting to the use of it, is no longer appropriate. It may also simply reflect the socioeconomic changes that the community went through, which may have eroded the knowledge and confidence in interpreting traditional indicators.

Yet in the process of the conversation, participants in the Chato Grande assessment also highlighted a range of indicators. The importance of environmental clues furthermore became apparent in the discussion of the flood of 2002. The community ignored warnings from outside sources, because floods were previously only experienced in conjunction with preceding heavy rains. As there were no rains, the warnings did not appear credible. The flood was triggered by heavy rains in the highlands.

Puno

The communities in Puno highlighted a broad number of indicators for the range of climatic risks they are exposed to (Table 16). While there is, however, a general confidence in forecasting wet or dry climate conditions or obtaining at least some warning of flood or frost conditions, there is low confidence in predicting hail events. Hail events are not understood and are strongly related to superstitious beliefs, being considered a form of punishment for the communities. This is reflected by indicators named by communities for hail events, which are predominantly linked to human behavior associated with guilt.

Table 16. Indicators Used by Communities to Predict Weather Extremes and Climatic Conditions

Wet/Dry Conditions

Category	Indicator	Type of Observation	Community												Example (an initial estimate of the validity and practicality as indicator is included. [likely/questionable/unlikely])
			Piura			Candile						Puno			
			Pq	CG	Loc	Cb	Ay	Anc	SM	CC	Ca	Ym			
	Temp	Felt temperature anomalies in the months preceding the rainy season	-	Y/-	-	-	-/-	-/-	-/-	-/-	-/-	-/-	-/-	Chato Grande (Piura): Exceptionally hot temperatures in November and December are signs of abundant rains in February and March. [likely] Locuto (Piura): When it is exceptionally hot and windy, this is indicative of a strong rainy season. [likely]	
	SST	Ocean temperatures in the months preceding the rainy season	Y/-	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.	Parachique (Piura): Exceptionally warm SSTs are indicative of abundant rains. [likely]	
	Wind	Direction, strength, months of occurrence	-	Y/Y	Y/-	-	Y/Y	-/-	-/-	-/-	Y/Y	Y/Y	Y/-	Chato Grande (Piura): Whirlwinds going from north to south in November and December indicate rain, while whirlwinds in September and October indicate drought. [likely] Locuto (Piura): Winds from south to north in October and November indicate a dry year. [likely] Aychuyo (Puno): Whirlwinds from the lake suggest that parts of the year will be rainy. [questionable] Ancaca, Candile, Cari Cari (Puno): Strong winds in July and August coming from Lake Titicaca indicate a rainy year. [questionable]	
	Fog, Mist	Appearance	-Y	-	-	-	Y/-	-/-	-/-	Y/Y	Y/Y	-	Y/-	Santa Maria, Aychuyo, Candile (Puno): Mist covering the entire community is seen as indicator of excessive rain and potential flood risk. Absence of mist in the hills is considered a sign of drought in the community of Santa Maria. [questionable]	
	Snow	Abundance	-	-	-	-	-/-	-/-	-/-	-/-	-/-	-/-	-/-	Yanamocco (Puno): Abundant, black clouds appears to be a short-term indicator and warning of strong precipitation. [likely]	
	Clouds	Appearance	Y/-	-	-	-	-/-	-/-	-/-	-/-	-/-	-/-	-Y	Locuto (Piura): Good visibility of the Milky Way indicates a good/rainy year [likely] Santa Maria: Small, brilliant, and close appearance of stars is seen as a sign of a rainy year. [likely]	
	Stars	Appearance, brightness	-	Y/-	Y/-	-	-Y?	-/-	-/-	Y/Y	Y/Y	-	-Y	Chato Grande (Piura): If the moon's limbs are pointing north it will be rainy; if they are facing south the year will be dry. [unlikely] Santa Maria/Aychuyo (Puno): If the new moon has a yellow color, this is considered a warning of excessive rain and flood risk. [unlikely]	
Physical	Moon	Appearance, color	-	Y/Y	-	Y/Y	-/-	Y/-	-/-	Y/Y	Y/Y	-	-/-		

Note: Pq = Parachique; CG = Chato Grande; Loc = Locuto; Cb = Cabuyal; Ay = Aychuyo; Anc = Ancaca; CC = Cari Cari; SM = Santa Maria; Ca = Candile; Ym = Yanamocco; Y = yes; n.a. = not applicable. The authors recommend that indicator examples classified as either "likely" or "questionable" are investigated further in their practical potential. Source: Authors' analysis.

Table 16. Indicators Used by Communities to Predict Weather Extremes and Climatic Conditions (continued)

Category	Indicator	Type of Observation	Community										Example (an initial estimate of the validity and practicality as indicator is included. [likely/questionable/unlikely])		
			Piura			Puno				Candile					
			Pq	CG	Loc	Cb	Ay	Anc	SM	CC	Ca	Ym			
Plant	Plant	Appearance, timing of growing stages	-	Y/-	-	Y/-	Y/-	Y/Y	-Y	Y/Y	Y/Y	Y/Y	Y/Y	Y/Y	Chato Grande (Piura): Delayed and abundant flowering of carob tree indicates a rainy year. [questionable] Cabuyal (Piura): The abundance of wild berries from October to December before the rainy season indicates the year will be good for production. The blooming of the guabo silvestre from July to August and abundant flowering in the lanche and arrayan trees are considered reliable signs of a good/rainy year. [questionable] Santa Maria (Puno): The appearance of bright green algae is a sign of abundant rains. [questionable] Santa Maria, Candile, Cari Cari (Puno): Early and/or abundant flowering of kanwa signals a rainy year. [questionable]
	Insects	Appearance, abundance, behavior	-	Y/-	Y/-	Y/-	-Y	-Y	-Y	-Y	?	-/-	-Y	-Y	Cabuyal (Piura): The abundance of red ants is a sign that water will be abundant. [questionable]
	Spiders	Appearance, behavior	-	-	-	-	-/-	-/-	Y/Y	Y/?	Y/-	-/-	-/-	-/-	Santa Maria/Cari Cari (Puno): Abundant spiders in the straw indicate rain, while spiderwebs in low part of the grasses signal dry conditions. Excessive numbers of spiderwebs are considered a sign of abundant rains in community of Cari Cari. [questionable]
	Snake	Appearance, abundance, calls	-	-	-	-	-/-	-/-	Y/-	Y/-	-/-	-/-	-/-	-/-	Appearance of black snakes between August and December signals a wet, rainy season. [questionable]
	Amphibians	Appearance, abundance, calls	-	-	-	Y/-	-Y	-Y	?	-/-	-/-	Y/?	-Y	-Y	Cabuyal (Piura): The constant croak of the toad before the rainy season indicates a good year. [questionable] Candile (Puno): If toads move to higher parts, this is taken as a sign of wet conditions. [questionable]
Biological	Fish	Abundance, behavior	-	-	-	-	-/-	-/-	-Y	-/-	-/-	-/-	-/-	-/-	Locuto (Piura): When the chilalo nests in the highest places, it will be a rainy year. When it nests near the river this indicates a dry year. While previously considered a relevant indicator, the community noted that this indicator has failed since 1998. [questionable]
	Birds	Appearance, flight pattern, nesting behavior	-	Y/Y	Y/Y	Y/Y	-Y?	Y/Y	Y/Y	Y/Y	Y/Y	Y/Y	Y/Y	Y/Y	Cabuyal (Piura): When herons leave the coastal areas and go to higher areas in Piura, this indicates a dry year. Puno: With the exception of Aychuyo, all the other five communities assessed in Puno highlight the egg-laying behavior of Leke Leke and color of the eggs as good indicator of drought. Four of these also use the bird's nesting behavior as an indicator of excessive rains and flood risk. [questionable]
	Mammals	Appearance, behavior	-	-	-	-	-Y	Y/Y	-Y	Y/Y	Y/Y	Y/-	-Y	-Y	Anccaca (Puno): If foxes have their offspring in the hills, this is seen as indicative of flood risks. [questionable]

Note: Pq = Parachique; CG = Chato Grande; Loc = Locuto; Cb = Cabuyal; Ay = Aychuyo; Acc = Anccaca; CC = Cari Cari; SM = Santa Maria; Ca = Candile; Ym = Yanamocco; Y = yes; n.a. = not applicable. The authors recommend that indicator examples classified as either "likely" or "questionable" are investigated further in their practical potential. Source: Authors' analysis.

Table 16. Indicators Used by Communities to Predict Weather Extremes and Climatic Conditions (continued)

Category	Indicator	Type of Observation	Community												Example (an initial estimate of the validity and practicality as indicator is included. [likely/questionable/unlikely])
			Pitira						Puno						
			Pq	CG	Loc	Cb	Ay	Anc	SM	CC	Ca	Ym			
Other	Religious Holidays	Link with physical observation	-	-	-	-	Y	Y	Y	Y	Y	Y	Y	Y	Religious holidays are used to reference points in time. Physical observations at the time of religious events are then taken as indicators of the characteristics of the upcoming rainy season. Holidays highlighted when observations concerning the quality of the rainy season are made include Concepcion, San Juan, San Andres, San Antonio, San Jose, San Pedro, and Santa Teresa. Whether there are relevant links between the observations and the characteristics of the rainy season needs to be evaluated for the various dates of the holidays. [questionable]
		Link with human behavior	-	-	-	-	-	-	-	Y	-	-	-	Y	Here, observations at religious holidays reflect superstitious beliefs of interlinkings between human behavior (good/bad) and the appearances and quality of the rainy season. Predictive relevance for climatic conditions can be ruled out, but the importance of such ingrained beliefs to communities needs to be considered. [unlikely]

Note: Pq = Parachique; CG = Chato Grande; Loc = Locuto; Cb = Cabuyal; Ay = Aychuyo; Acc = Anccaca; CC = Cari Cari; SM = Santa Maria; Ca = Candile; Ym = Yanamocco; Y = yes; n.a. = not applicable. The authors recommend that indicator examples classified as either "likely" or "questionable" are investigated further in their practical potential.
Source: Authors' analysis.

Table 16. Indicators Used by Communities to Predict Weather Extremes and Climatic Conditions (continued)
Hail

Category	Type of Indicator	Observation	Community					Example (an initial assessment of the validity and practicality as indicator is included [likely/questionable/unlikely])
			Ay	SM	Anc	Ca	CC	
Physical	Clouds	Dark clouds	Y	Y	Y	Y	Y	Short-term indicator shared by all the communities. [likely]
	Sun/temperature	Felt temperature	Y	Y	Y	Y	Y	Short-term indicator shared by all communities, except Cari Cari. [questionable]
	Wind	Strength or direction		Y	Y	Y	Y	Location-specific short-term indicator. Santa Maria observes southern winds from the hills in advance of hailstorms, while the communities Ancacca, Candile, Yanamocco highlight very strong winds, which in the case of the latter two communities come from a western direction. [questionable]
Animals (selection)	Spider	Behavior	Y	Y	-	-	-	The behavior of spiders (collection of white eggs) is considered an indicator of hail events for community members in Aychuyo and Santa Maria. [unlikely]
	Ant	Behavior	Y	-	-	-	Y	The behavior of ants (collection of white eggs) is considered an indicator of hail in Aychuyo and Yanamocco. [unlikely]
	Bird	Behavior	Y	Y	-	Y	Y	The nesting behavior of leke leke (nests located in the hills; dark eggs) is taken as a sign of a year with hail events. [unlikely]
Plants (selection)	Fox	Behavior	-	-	Y	-	-	The community members in Ancacca consider long howls a sign a bad year with hail events. [unlikely]
	Algae	Physical appearance	Y	-	Y	-	Y	The appearance of algae is taken a sign for hail conditions in Aychuyo, Ancacca, and Cari Cari. [unlikely]
	Potato	Physical appearance	Y	-	-	Y	Y	If the flower appears larger than usual or leaves are broken (Candile) or leaves appear fragile (Aychuyo, Cari Cari, Yanamocco) this is taken as sign for hail conditions. [unlikely]
Other	Various	Timing of bloom	-	Y	Y	-	-	In particular, community members of Ancacca consider the timing of the blooming of various local plants such as tola an indication of whether the climatic conditions in the year are conducive to hail or not. [unlikely]
	Women, household	Abortion	Y	Y	Y	Y	Y	Widespread belief in communities that abortions will be "punished" with hailstorms. [unlikely]
	Children	Punishment, crying, playing naked	Y	-	Y	Y	Y	Behavior deemed negative toward or by children in the communities is considered linked to the subsequent occurrence of hail. [unlikely]
Dreams	Community Members	Fights	-	-	-	Y	-	Conflict within the community promotes punishment through hail events. [unlikely]
	Women	Appearance after birth	Y	-	-	-	-	Women who give birth should not show themselves in the sun to avoid negative influence on the climate. The blood of birth is considered an impurity and, therefore, the woman has to conceal herself. [unlikely]
	Dream	Negative images in dream (e.g., fights)	-	Y	Y	Y	Y	Negative images of human behavior transported in dreams are associated with subsequent hail events. [unlikely]

Note: Pq = Parachique; CG = Chato Grande; Loc = Locuto; Cb = Cabuyal; Ay = Aychuyo; Acc = Ancacca; CC = Cari Cari; SM = Santa Maria; Ca = Candile; Ym = Yanamocco; Y = yes; n.a. = not applicable. The authors recommend that indicator examples classified as either "likely" or "questionable" are investigated further in their practical potential. Source: Authors' analysis.

Table 16. Indicators Used by Communities to Predict Weather Extremes and Climatic Conditions (continued)

Frost

Category	Type of Indicator	Observation	Community						Example (an initial assessment of the validity and practicality as indicator is included [likely/questionable/unlikely])
			Ay	SM	Anc	Ca	CC	Ya	
Physical (Selection)	Stars	Appearance	Y	Y	Y	Y	Y	Y	Appearance of the Cotto Constellation in January and February is a sign of whether frosts are likely in the season. [questionable]
	Wind	Direction	Y	Y	-	-	Y	Y	Winds blowing from the mountains are considered a sign of impending frost conditions, particularly in the communities of Aychuyo and Yanamocco. [questionable]
Animals (Selection)	Bird	Behavior	Y	Y	-	-	Y	Y	The location of the nests of leke leke is considered an indicator of the characteristics of the rainy season. Location of nests in the lowland and dark eggs are taken as signs of a dry year, as well as increased likelihood of frosts. [questionable]
	Various arthropods, insects, and amphibians	Behavior	Y	Y	-	-	Y	Y	Various forms of behavior are linked to a range of animals that are associated with increased likelihood of frost conditions. These, however, are listed only in individual communities and did not receive broad support. [unlikely]
Biological (Selection)	Plants	Flowering, fructification	Y	Y	-	-	Y	Y	No or delayed flower building and fructification in a variety of local plants is taken as a sign of climate conditions conducive to frost. Plants of interest include karihua, tola, and sankayo. [questionable]
Other	Religious Holidays and Specific Dates of Observance	Physical characteristics and specific dates	Y	Y	-	-	Y	Y	Various holidays and dates are used as reference points for the observations of atmospheric conditions as indicators of climate conditions conducive to droughts and frost. The absence of clouds at specific holidays is widely considered a sign that frosts are likely. Highlighted holidays include Nuestra Señora de la Paz (January 6), San Sebastian/Candelaria, and San Andres. Specific dates highlighted for observation include June 23 and 29, August 1–3, 15, and 31. [questionable]

Note: Pq = Parachique; CG = Chato Grande; Loc = Locuto; Cb = Cabuyal; Ay = Aychuyo; Acc = Anccaca; CC = Cari Cari; SM = Santa Maria; Ca = Candile; Ym = Yanamocco; Y = yes; n.a. = not applicable. The authors recommend that indicator examples classified as either "likely" or "questionable" are investigated further in their practical potential.

Source: Authors' analysis.

Gauging the Role and Validity of Indicators

Piura. The physical indicators highlighted by the communities of Piura warrant further investigation concerning their reliability and potential to complement scientific forecasts.

The association of exceptionally warm SSTs in the months preceding the rainy season represents a valid observation of the evolution of local El Niño conditions, which are associated with the cold, nutrient-rich waters with warm ocean (correntada) currents. Exceptional land-surface temperatures are also a consequence of local El Niño conditions in the months preceding and during the rainy season.

As local El Niño conditions are associated with the prolongation and intensification of the rainy season, early precipitation events generally represent a valid signal for the communities to prepare for an intense rainy season as well as heightened flood risk.

While intense precipitation events represent a warning of possible flooding in the Piura River basin, the absence of rainfall in the lower Piura River basin does not necessarily preclude a flood risk. The flooding of Chato Grande in 2002, where warnings were largely ignored due to the absence of intense rainfall, illustrates the need to strengthen the capacity of the communities in the river basin to relate their local observation with more regional forecasts. This applies also to the dissemination approach of scientific climate forecasts and early warning, which must recognize the important influence of local observations.

Wind direction was also highlighted by communities in Piura as an important indicator of wet or dry conditions. This appears to be a valid indicator for the coastal areas of northwestern Peru, which also has been noted in coastal Ecuador (Cornejo, personal communication). During El Niño years, the southward displacement of the Intertropical Convergence Zone would introduce a circulation anomaly, which may be observable in the prevalent wind directions or

cloud movements. To gain further confidence in this indicator, the reliability of these observations would need to be tested with a high-resolution time series on wind patterns, which was not available in time for this report.

With the exception of Parachique, all communities in Piura highlighted a range of biological indicators. Given that changes in temperature, humidity, and radiation affect the physiology of plants, the semiperiodic fluctuation between dry (cool) and wet (hot) conditions due to ENSO makes it likely that these local plants may provide valuable clues to the characteristics of the rainy season. Similarly, animals can function as indicators if their appearance or behavior is closely tied to temperature or moisture profiles associated with El Niño events or dependent on plant species that are linked to these climatic conditions. Longer-term observations would be required to identify the lead times and reliability of local plant and animal species as indicators and to differentiate these indicators from spurious correlations and false knowledge.

Puno. Research has shown that the practice of indigenous Andean communities of forecasting the quality of the rainy season according to the brightness of the Pleiades is based on sound observation and yields equal or better predictive capabilities than some climate forecasts for the region (Orlove et al. 2000). In the assessments, the communities placed their highest confidence in predicting dry or wet conditions, while they only had low confidence in forecasting hail conditions.

For forecasting the rainy season and associated flood risk, the communities highlighted a range of physical and biological indicators. Aside from the observance of stars, the observation of wind direction and strength is given considerable attention. Easterly winds are likely to be a valid physical indicator of a strong rainy season and heightened flood risk, as highlighted by the community of Candile. It has been shown that precipitation events during the austral summer months

are associated with mid-tropospheric winds with a large easterly component, which allows for the transport of moisture from the Amazonian lowlands into the altiplano (Garraud et al. 2003). Increases in these easterly flows would cause increases in precipitation.

Another indicator of wet conditions is ground fog, which is indicative of high soil-moisture content and reflects on the atmospheric moisture content of the region. The frequent observations could hence provide a valid near-term indicator of conditions conducive to intense precipitation or lakeside flooding, as indicated by the communities of Santa Maria, Aychuyo, and Candile.

The observation of clear skies is considered an indicator of dry conditions favoring the occurrence of frosts. Given the high altitude of the altiplano, this is likely to be a valid inference if clear skies are persistent during the night and the day (Seimon 2006). This would be indicative of pronounced atmospheric moisture deficit, which would favor strong cooling overnight due to the absence of insulating cumulus clouds.

Another likely valid short-term indicator of frosts in the Titicaca region is southwest winds. Southwest winds would be downslope winds descending from cordilleras in the west, displacing moist air from the Amazonian basin (Seimon 2006). If this anomalous wind direction is sustained, it would lead to a drying of the atmosphere (boundary layer) and hence favor conditions that promote overnight cooling (as described above).

Concerning the physical data, meteorological observations in communication with the communities would help in providing further detail on the lead times and reliability of these observations. As already pointed out for the Piura assessments, this could also help to relate micro- and mesoscale observations with larger-scale regional forecasts. The absence of accessible meteorological data with high temporal resolution and longer-term records represents a major obstacle in validating the observations from the communities.

Nevertheless, systematically recording the observations of the communities and relating them to the occurrence of climatic extremes and scientific forecasts could provide important clues in relating regional scientific forecasts and local knowledge.

Perceptions of Changing Climate Conditions

Assessments in northwestern Peru and the altiplano region of southern Peru independently revealed a concern among the communities about observed climatic and environmental changes. The perceptions described below are the outcomes of discussions with community representatives during the synthesis workshops in Piura (communities from Piura) and Lima (communities from Puno).

Piura: Perceived Climatic Changes in Northwestern Peru

In northwestern Peru, a major concern is the perceived increase in daily maximum temperatures, while nighttime temperatures have become colder. While dry conditions are the norm in this region of Peru, the coastal communities felt that the drought conditions they have been exposed to over the recent years were unusual in their duration and pervasiveness.

Puno: Perceived Climatic Changes in the Peruvian Altiplano

In the altiplano region, communities are also experiencing increases in daily maximum temperatures, saying they need to wear more sun-protective clothes such as hats than they used to when working in the field. Nighttime temperatures also appear to have become colder, falling more often below freezing. This adversely affects important crops such as sweet potatoes during the planting season.

A major concern in the altiplano is the perceived change in the timing and magnitude of precipitation during the rainy season. Rains started around mid-December 15 to 20 years ago and used to peak in intensity around February and March before fading out in April, which is closely resembles the historical data record for Puno from the 1930s to 70s (see Fig. 3) While the beginning of the rainy season does not seem to have changed, communities repeatedly noted that no rains or only very little rainfall occurs during February and March. Instead, more frequent cold spells are observed. This is particularly detrimental to the potato harvest, as during this time, the plant is flowering and most vulnerable. Heavy rains are now observed in April, when the potatoes would usually be harvested.

Another worry is the decrease in water flows. The community representatives felt if this trend continues, they will have little or no water within the next one to two decades.

Are These Perceptions True?

A continuous challenge in validating community perceptions about climatic changes is the access to relevant data. In the case of Peru, the raw data may have been collected by the meteorological service (SENAMHI), but has not been processed aside from the main meteorological stations due to staffing limitation (personal communication with representatives from SENAMHI). Data can only be provided through costly formal requests, which are in part needed for covering staffing costs, but are also a common practice in many developing countries.

Besides accessibility, the quality of data may be a constraining factor. Short or discontinuous climatic records make it difficult to detect and validate trends. The lack of access to quality meteorological data constrains research efforts and may lead to a late detection of climatic signals, which ultimately have practical implications for climate-sensitive development processes, such as agriculture.

Another challenge is to correctly relate the scale of observations by the communities with the resolution of the data and presentation of data. Monthly or seasonal averages compiled may smooth out and hence mask signals, which are recognized by farmers and other individuals living closely with the environment.

The discussion presented below does not pretend to be comprehensive; rather it is an attempt to obtain an initial insight into whether some of the community perceptions are reflected in meteorological observations. Emphasis is placed on the main climatic parameters; i.e., average, maximum, and minimum temperatures and precipitation. Monthly and seasonal summaries are usually presented. Where possible, additional analysis is included or reference to published research is made.


The analysis below should be considered a start in guiding further analysis. In order to detect climatic signals at the community level, it appears to be valuable to include more systematic observations from farmers.

PIURA


Figure 12 shows a warming trend in the daily maximum temperature for the months of January, February, and March for Esperanza in the coastal region of northwestern Peru. The discontinuous record initially also shows increases in the daily minimum temperatures for the same months. Following data gaps more recent values show a steep increase in maximum temperatures and a steep drop in minimum temperatures. Whether this reflects the recent perceptions of the communities that daytime temperatures are getting warmer while nighttime temperatures are getting cooler or is simply the result of other factors, such as changes in instrumentation is unclear. Given the discontinuous record, this interpretation has to be viewed with caution. It also contradicts the general global trend of decreasing diurnal temperature ranges (e.g., IPCC 2001).

Figure 12. Average Monthly (A) Maximum and (B) Minimum Temperatures for Esperanza, Northwestern Coastal Peru

A



B



Note: For the monthly averages of the hot months of January, February and March an increase in the (a) maximum temperatures can be observed from 1972 to 1996 can be observed. For the continuous data period from 1972–1993 (b) minimum temperatures also appear to be increasing. It is unclear, based on the limited measurement points, whether the steep increase in maximum temperatures and drop in minimum temperatures after the data gap is a result of change in instrumentation or reflects the more recent perceptions of the communities.

Source: CONAM.

For Huancabamba in alto Piura, positive trends or no trends in daily maximum temperatures are observed for the months of January to March (Figure 13). While the maximum temperatures for January appear to show an increase over time, the minimum temperatures exhibit a slight decrease over time, and the opposite appears to be the case for the months of March. The perceptions of the communities are not clearly reflected in the data. In general, the difficulty to obtain recent and continuous observational data records, constrains the ability to draw clear conclusions.

PUNO

Changes in the temperature were also a concern in the altiplano. The community representatives noted that it appears to get hotter during the day, while colder at night.

It was not possible to obtain long-term temperature records for the department of Puno for assessing the validity of this observation. Instead, long-term records obtained for Cusco were taken as a proxy. It can be expected that observed trends in maximum temperatures have the same sign. Due to the geographical distance the analysis of minimum temperature trends have to be viewed with greater caution, as microclimatic factors play a greater role.

Figure 14 shows that over a period of more than 40 years a significant warming trend can be observed in minimum and average temperatures, but not in the maximum temperature.

The temperature data from Cusco does not exactly match the observations in the communities. While a warming trend can be detected, it is observed for minimum and average temperature averages and not for maximum temperatures. The temperature data, however, reflects also the general observation that the warming trends observed across the globe tend to be more pronounced in minimum temperatures than maximum temperatures (IPCC 2001). Research

conducted in the Bolivian and Peruvian parts of the Andes generally confirms though that the region is getting warmer; a warming trend can be detected at all elevations on the Pacific side of the Andes. General temperature increases in the range of 0.15 to 0.39 °C per decade have been observed over recent decades within the region (see Vuille and Bradley 2000, Vuille et al 2003, Diaz et al. 2003).


While microclimatic differences cannot be ruled out between Cusco temperature record presented here and the region around Puno, it is likely that the general warming trend is likely to be similar. What could therefore lead to the vivid perception by community representatives that it is particular the maximum temperatures, which have increased?

Time records suggest that there is an increase in the intensity of solar radiation received in the altiplano over time and the also appears to be an extension in the number of sun hours per day (Seimon, personal communication). Given that the community perceptions are based on felt increases in temperatures, it may be that their response rather reflects the increased intensity of the sun during the day rather than changes in the daily maximum temperature.


The perception of the communities that there are changes in precipitation characteristics of the altiplano is reflected in precipitation data in Figure 15. Precipitation totals for the months of January, February, March, and April averaged over 30 years and an area of 0.5 x 0.5 degrees, suggests that there is a shift from the maximum precipitation amount occurring in January toward March over time. While these are broad averages, the direction of change is in line with the community perception that there is a shift in the general precipitation pattern. If this can be confirmed on finer temporal and spatial scales, this would have implications for the time of planting cycles and may explain some of the irritation of the communities with the reliability of environmental indicators.

Figure 13. Average Monthly (a) Maximum and (b) Minimum Temperatures for Huancabamba, Sierras, Upper Piura Watershed

A



B




Note: Monthly averaged (a) maximum and (b) minimum temperatures are displayed. Positive trends or no trends in daily maximum are observed for the months of January to March. While the maximum temperatures for January appear to show an increase over time, the minimum temperatures exhibit a slight decrease over time, and the opposite appears to be the case for the months of March. The perceptions of the communities are not clearly reflected.


Source: CONAM.

Figure 14. Long-term Temperature Record for (a) Minimum, (b) Maximum, and (c) Average Temperature in Cusco, Peru

A




B



Note: While no trend in the maximum temperature can be detected, a warming trend in minimum and average temperatures can be detected (processed data courtesy A. Seimon).


Figure 14. Long-term Temperature Record for (a) Minimum, (b) Maximum, and (c) Average Temperature in Cusco, Peru (continued)



Note: While no trend in the maximum temperature can be detected, a warming trend in minimum and average temperatures can be detected (processed data courtesy A. Seimon).

While further detail is needed to explore the interaction with interannual and interdecadal climate variability, the perceptions of changing climatic conditions and concern for water resources voiced by the communities should be taken serious. Aside from scientific trends, adaptation responses are already being reported. Halloy et al. (2005) have documented an upward shift in the area of cultivation in some parts of the altiplano. A shift in the precipitation patterns in conjunction with the ongoing glacial melt would have considerable implications for agricultural activities in the altiplano.

Figure 15. Precipitation Averaged over 30-year Periods for January to April for the Peruvian Altiplano Region



Note: The data suggest a possible shift in timing of the rainy season over time.
Data source: Climatic Research Unit, provided by CIP.

3 Conclusions and Recommendations

The assessments illustrate that strengthening the adaptive capacity to climatic shocks is vital to the sustainable development of the coastal and highland regions of Peru. A major challenge for the communities in Piura and Puno is to protect critical assets in order to avoid a poverty spiral. The important aspect here to consider is that development activities have to be conducted with a full appreciation of the risk envelope of the communities.

Both in northwestern Peru and in the altiplano, managing risks across the entire spectrum of climate variability becomes more important in light of climate change. In Piura, this means better managing the fluctuations between dry and wet climatic conditions. El Niño events have to be managed not only in terms of flood risks, but also viewed as an opportunity to livelihoods due to the above normal water supply, revitalized vegetation, and so on. These positive effects for natural-resource-dependent livelihoods, can only be utilized, however, if these livelihoods are better (i) buffered against the destruction of infrastructure, disease outbreaks, and other impacts associated with the climatic extremes of strong El Niño events, and (ii) equipped to utilize the subsequently improved environmental conditions through access to markets and credit. The semiperiodic occurrence of ENSO events and the fairly clear-cut risk exposure provides a good planning framework for strengthening the resilience of communities to climatic risks.

In Puno, managing climatic risks means strengthening resilience to a broad spectrum of climatic extremes

that can occur in short sequence. These risks make it very difficult for communities to accumulate assets or regain them once they are lost in a disaster event. There is a need to improve the yield of existing natural-resource-based livelihood practices, while also providing the opportunity for diversifying into income-generating activities that are more climate-resilient. The multihazard environment makes access to credit and insurance mechanisms paramount to minimize the impact of climate-related shocks on households. Here, innovative initiatives suited to this high-risk environment are needed.

The report represents only a point-in-time investigation of the perceptions of communities concerning their exposure to climatic risks and capacity to adapt and cope. The exercise has shown, however, a strong willingness and interest by the communities to better manage environmental risks. But the communities are certainly aware of their limitations, and in order to improve their situation outside support is needed in terms of information that supports the identification of risk factors (especially helping to clarify causal and spatial relationships), informs them about adaptive measures, and provides an enabling environment for creating more climate-resilient livelihoods. The sections below highlight a range of interventions that can be undertaken at the national, department, and community level and are likely to reduce vulnerabilities. Only general cross-cutting issues that apply to Piura and/or Puno communities are identified, while community-specific interventions are listed in Appendix D.

Use of Climate Information

The assessments have illustrated that building a bridge between external information and internal decision-making structures and knowledge are important in the adaptive management of climate risks.

El Niño forecasts are predominantly recognized by Piura communities and not communities in Puno, where the effects of El Niño are masked by the multihazard environment. Even in Piura, however, the recognition of El Niño events differs considerably from scientific classification. Only the strong El Niño events of 1982–83 and 1997–98 were picked up by the communities, while more moderate events in between this time frame were usually not recognized by the communities as El Niño events. Furthermore, La Niña is not a concept the communities tend to be familiar with. This has to be reflected in the dissemination of early warnings of ENSO events.

For the communities of Puno, the climate-risk profile makes more holistic access to weather and climate forecasts more important than warnings of particular El Niño events. In both cases, it is important the information formats fit into the local knowledge and decision-making structures.

The review of local indicators on climate and environmental conditions suggest that they are still of considerable importance to the communities, although the knowledge of these indicators appears often rudimentary or confined to a smaller group.

Through ENFEN, the government of Peru has a promising network of institutions engaged in information collection and forecasting. The main constraints that form the perspective of communities concerning the use of information based on these forecast capabilities of El Niño conditions appear to be lack of trust toward some of the information providers and the dissemination method.

In Puno, efforts need to be more focused on improving the understanding of climate risks in their spatial dimension at the community level and providing climate information that is more directly linked to the production systems of the altiplano. A major constraint is the dissemination format, since such a format needs to take into account the low literacy levels of the communities.

Activities focused on improving the integration of climate information into the decision-making processes at the community level could include:

- (i) Trust-building exercises between information providers and communities through local meetings;
- (ii) Development (staffing/training) of multiple extension services that link information on climate risks with practical guidance on natural resource management, agricultural and livestock practices, health and hygiene practices, and disaster prevention and allow for a two-way dialogue process with the communities;
- (iii) Collection and assessment of local knowledge of environmental indicators building on existing information collection structures by the General Directorate for Agrometeorology (GDA) within SENAMHI and other institutional structures; and
- (iv) Review and reform of the consensus-finding process on ENSO conditions to allow for more timely dissemination of forecasts based on a more participatory dissemination process, enabling a more detailed communication of uncertainties and improvement.

Research

The focus of the recommendations herein is on applied research that aims to relate the scientific analysis of the physical and economic environment to the practical needs of the communities.

Environmental Risks and Change

The discussions with the communities have shown that there is a widespread perception of change. Some of these changes could be confirmed in data records. There is, however, the need for more in-depth investigation of climatic data and environmental signs at scales that are most relevant to the livelihood frameworks of the communities. It is not only important to detect whether there are changes in the absolute monthly or annual values over time, but to what extent changes are occurring at temporal scales that are directly relevant to the livelihoods of the communities (e.g., changes in the characteristics of the rainy season).

Given the sensitivity of Piura and Puno to changes in water resources, it is important that a continuous monitoring of relevant climatic parameters is ensured in the regions. Availability and access to data is essential for the early detection of climate signals in these regions, as this may have profound implications for the development.

In light of climate change, detailed profiles of current and projected climatic conditions for the various departments of Peru, and understanding how these profiles relate to the prevalent livelihoods, is becoming ever more important. CONAM has embarked on evaluating climate change in various departments, conducting basin-wide assessments under, for example, the ProClim program. This is a promising development; however, based on the consultations for this report, access to climate data is difficult for external researchers and often comes at a considerable cost. There is a need to make climate data more easily accessible to stimulate research activities on Peru, given its high vulnerability to climate variability and change.

Initiatives should particularly be focused on promoting research and facilitating exchange of methodologies and knowledge concerned with information on (i) climate baseline, (ii) climatic trends at scales relevant to development intervention, and (iii) refinement of detailed future climate scenarios.

There is also a need to strengthen the exchange between Spanish- and English-speaking research activities by supporting partnerships and translating important knowledge products.

Climate Forecasts and Local Indicators

The assessments in Piura and Puno illustrated a range of indicators that could help to relate forecasts to more-local climatic conditions. Given the importance of environmental signals to the acceptance of climate forecasts, further efforts should include a systematic assessment of the validity of promising biophysical indicators.

Agriculture and Livestock Management and Market Research

Research initiatives focused on promoting climate-resilient agricultural practices, pest management, and the protection of livestock needs to be promoted to improve the portfolio of practical options to reduce the impact of environmental hazards on the assets of the households. This should be connected with exploring market opportunities, such as identifying niche markets for (native) crops well-suited for the climatic conditions of the altiplano.

Incentive Systems

Communities in Piura and Puno both noted the absence of credit systems adequate for their specific needs. In Puno, organizations like CIRNMA have been successfully piloting credit schemes; however, there is a need to understand how such credit schemes and insurance mechanisms could be upscaled and sustained.

The effects of environmental degradation on households were also recognized in the assessments. There is a need to provide communities with options and incentives to reduce local environmental impacts, which exacerbate risk exposure. Creating, for example, incentives for

afforestation in the upper Piura River basin would not only help to counteract erosion processes in communities in Cabuyal, but also help to reduce the silting of the riverbed in lower Piura and, thereby, flood risks. In this context it may be worthwhile to explore potential revenue streams to communities created through the emergence of the carbon market for afforestation activities.

Tracking Progress

Given the advances in climate information and natural resource management practices, there is the opportunity to develop a comprehensive risk management strategy for communities. In order to assess the success of such efforts, however, it would be crucial to track progress over time. This should include monitoring the uptake of external information and risk management strategies in targeted communities. By tracking household assets and climate hazards over time, this should provide insights into how successful current risk management options are in improving the resilience of communities. If successful, households should be able to increase their asset base over time while facing the same or increasing hazard exposure. This needs to be validated through long-term monitoring efforts.

These systems could be designed in such a way that they can also function as bottom-up early warning systems, which trigger emergency response mechanisms when household assets fall below critical thresholds. Such mechanisms have, for example, been successfully developed in Kenya with financing from the World Bank.

Community Action on Climate Risk Management

The above section has already highlighted the need to tailor climate information more specifically to community-specific needs and improve communication structures that allow for feedback mechanisms and

integration of local knowledge. The adaptive capacity of communities needs to be strengthened through a combination of awareness-raising activities and targeted investments. A strong emphasis on awareness of risks and appropriate adaptation measures is needed in order to empower communities to make decisions on how they can address these risks and create ownership of local-level actions.

Awareness Building

While communities are aware of their vulnerability and some constraining factors to adaptive actions, the assessments have also shown that there often is still a limited knowledge of the mechanisms that increase vulnerabilities to a particular hazard, lack of understanding of spatial extent of disaster risk and linkages between environmental degradation and climate-related vulnerabilities, and a need for improved knowledge of adaptation options. Furthermore, there is also a lack of understanding of how changes outside their immediate environment could affect a community.

In Piura, Chato Grande serves as a case in point. By relating flooding conditions solely to the local presentation of rain, the community ignored warning of a flood, which was triggered elsewhere in the Piura River basin. In Puno, the communities lack, for example, an understanding of the mechanisms that trigger hail and hence link its occurrence to superstitious beliefs. When the communities engaged with CIRNMA in a dialogue on and spatial mapping of disaster risk, this sparked an increase in understanding the evolution of hazards and identifying areas at risk.

Activities focused on building community-level awareness and knowledge include

- (i) spatial mapping of disaster-prone areas using a participatory process that provides a learning environment;

- (ii) training on natural resource, agricultural, and livestock management techniques to mitigate the impact of climate risk and associated effects, and
- (iii) Initiation and strengthening of dialogue processes to strengthen knowledge exchange of adaptation and coping practices and promote community-owned solutions to risk management strategies that require intercommunity collaboration; e.g., community-focused flood early-warning systems and river-basin environmental management.

Community-Level Investments

It is suggested that awareness-raising efforts in the communities are coupled with concrete investments, which place an emphasis on community-centered microprojects. With the focus on risk management, communities could go through a modified community-driven development process aimed at creating ownership and commitment toward risk-mitigating measures.

Microprojects could be clustered around the following themes:

- (i) **Information Access/Infrastructure.** Microprojects under this category are focused on investments that improve communication infrastructure to better access information on climate risks, adaptation measures, and market information. Various dissemination tools such as mobile phones, radios, and computers could be tested with the communities based on community preferences and technical feasibility. This component would be coupled with training or education activities.
- (ii) **Livelihood Enhancement and Diversification.** Microprojects are focused on enhancing the resilience of current livelihood practices to climate risks, including, for example, promotion of technical interventions, application of intercropping and soil conservation techniques, and introduction of more resilient crop varieties, but also providing training

in alternative and less climate-dependent income sources.

- (iii) **Human Health.** Vector- and water-borne diseases have been closely linked to ENSO and are further affected by increased climatic variability, as evidenced in the repeated outbreaks of malaria (coastal Peru) and cholera. In the altiplano, the extreme fluctuations in temperature and inadequate protection of households constitute a major concern. Microprojects will focus on investments in hygiene and protection that minimize exposure to this risk. This also should be coupled with structural investments (see vii).
- (iv) **Livestock Health.** Investments aimed at improving livestock health by minimizing exposure and improving monitoring and treatment options are more urgently needed in the face of climate change.
- (v) **Environmental Rehabilitation.** Environmental degradation is a shared concern among communities in Piura and Puno, and training and targeted investments in environmental rehabilitation and natural resource management are needed.
- (vi) **Credit and Microinsurance:** Microprojects may include the piloting of community-focused, small-scale credit and insurance mechanisms to facilitate the adoption of new agricultural practices and other livelihood strategies.
- (vii) **Structural Investments:** This component would place an emphasis on strengthening the ability of communities to be prepared for increased climate variability and manage across the entire spectrum of climate conditions. This includes, for example, investments in water harvesting and storage technologies and erosion-control measures. Particularly in the altiplano, investments would also focus on reinforcing and improving the insulation of houses against extreme temperature fluctuations and providing low-cost options for the construction of shelters to better protect livestock and enhance productivity. Construction of safe storage facilities for seeds and forage represents another promising

investment for reducing the vulnerabilities of communities.

Structural investments would also focus on rehabilitating and training communities in the management of ancient structures or testing simple interventions based on transfer of knowledge from other regions, which have been shown to improve water use or reduce vulnerability to the extreme climate variability. In the altiplano, such efforts may include rehabilitation and training in the maintenance of waru warus (raised fields). In northwestern Peru, this may entail testing the suitability of albarradas (u-shaped detention ponds, pre-Inca technology that is still used in coastal Ecuador to capture excessive run-off and recharge aquifers) or sand dams, which can help hold water longer into the dry season.

Enabling Frameworks: Institutional Coordination and Support

The assessments have highlighted a range of initiatives, which are currently undertaken to reduce risks at the community level. However, most of these activities are confined to particular communities and there is a need to build a more comprehensive risk management framework.

There is a need for promoting partnerships at various levels, which lead to a more coordinate approach to managing climate-related risks. The workshops in Peru, which brought together community representatives and institutional representatives, illustrated how such meetings can break down barriers and provide learning opportunities between the various stakeholders. The communities from Puno (altiplano region) expressed the desire to continue this dialogue. Institutions may help to facilitate such meetings or use them to create awareness about their work. There is currently a rather weak link with municipalities resulting in a feeling of exclusion by the communities; strengthening these linkages through outreach events or by other means would help to increase awareness of options in the

face of disasters. The goal should be working with and expanding on the existing structures and integrating external partners where appropriate.

External assistance to communities focused on coping with a specific disaster should focus on solutions that also reduce vulnerabilities to future hazard exposure. Communities in northern Peru and the altiplano repeatedly noted the inadequacy of building design and infrastructure for the prevalent risk profiles. The destruction of school buildings or the lack of insulation to protect them from extreme temperatures had cascading effects on the development prospects of the communities by affecting the educational opportunities of the children. Therefore, incorporating risk profiling and mapping and building safety standards needs to be promoted when engaging with the communities in reconstruction efforts. Solutions will need to go beyond the immediate alleviation of the suffering in the face of extreme events by also focusing on reducing the likelihood of recurrence of such suffering. The emerging signs of climate change make the focus on preventive and adaptive measures to climate risks in Peru ever more urgent.

Peru's growing emphasis on disaster risk prevention and adaptation to climate change needs to be supported and developed. Given the high prevalence of natural disaster and the increasingly visible effects of climate change, risk management efforts have to be understood as an integral part of development processes and not as an add-on. Community-focused microprojects, such as the ones outlined above, represent the opportunity to pilot and then upscale interventions aimed at reducing the vulnerabilities of the poor. Such local efforts also need to be better linked to the relevant national and sub-regional support structures. While there is currently a range of governmental, non-governmental and scientific expertise available, initiatives are often limited in their scope and reach. While some communities have access to several organizations, others receive little or no support. It remains a challenge to build a broader enabling framework that helps to disseminate and

replicate positive experiences in managing climatic risks and promoting development in the region.

Aside from an emphasis on reducing the existing adaptation deficit to currently observed climate variability and change, institutional efforts need to continue strategic planning efforts for managing the medium- to long-term implications of climate change. This means stimulating research and knowledge sharing

in basic climate science for Peru, as well as targeting applied research into risk management options for vulnerable development sectors. In conjunction with developing the knowledge basis, efforts need to be made to identify processes in policies and education that need to be initiated now to ensure that future Peruvians have the option to pursue livelihood strategies that are less vulnerable to the impacts of climate change.

References

- AfDB (African Development Bank, ADB (Asian Development Bank), DFID (Department for International Development, U.K.), BMZ (Federal Ministry for Economic Cooperation and Development, Germany), DGIS (Ministry of Foreign Affairs—Development Cooperation, The Netherlands), OECD (Organisation for Economic Cooperation and Development), UNDP (United Nations Development Programme), UNEP (United Nations Environment Program), and the World Bank. 2003. *Poverty and Climate Change: Reducing the Vulnerability of the Poor through Adaptation*. Interagency report, F. Sperling, ed. Washington, DC. 43 pp.
- Burton, I., and M. van Aalst. 2004. “Look Before You Leap: A Risk Management Approach for Incorporating Climate Change Adaptation into World Bank Operations.” Environment Department Paper. World Bank, Washington, D.C.
- Emanuel, K. 2005. “Increasing Destructiveness of Tropical Cyclones over the Past 30 Years”. *Nature* 436, 686-688.
- CAF (Corporacion Andina de Fomento). 2000. *Las lecciones de El Niño. Memorias del Fenomeno El Niño 1997–1998 Retos y propuestas para la region andina*. Volumen V: Peru. CAF Publication. Caracas, Venezuela.
- Charveriat, C. 2000. *Natural Disasters in Latin America and the Caribbean: An Overview of Risk*. Washington, DC: Inter-American Development Bank.
- Francou, B., M. Vuille, P. Wagnon, J. Mendoza, and J.-E. Sicart. 2003. “Tropical Climate Change Recorded by a Glacier in the Central Andes during the Last Decades of the 20th Century: Chacaltaya, Bolivia, 16° S.” *Journal of Geophysical Research Letters*, Vol. 108: D5, 4154.
- Francou, B., P. Ribstein, P. Wagnon, E. Ramirez, and B. Pouyaud. 2005. “Glaciers of the Tropical Andes: Indicators of the Global Climate Variability.” In: *Global Change and Mountain Regions: A State of Knowledge Overview*. U. Huber, K. M. Harald, and M. A. Reasoner, eds. New York: Springer: pp. 197–204.
- Garcia-Godos, J., D. McNeill, K. O’Brien, and L. O. Naess. 2004. “Seasonal Climate Forecasts and Community Responses: Summary of Experiences in Peru and Ecuador.” Consultancy Report submitted to the Climate Change Team, World Bank.
- Garreaud, R., and P. Aceituno. 2001. “Interannual rainfall variability over the South American Altiplano.” *Journal of Climatology*, 14: 2779-2789.
- Garreaud, R., M. Vuille, and A. Clement. 2003. “The Climate of the Altiplano: Observed Current Conditions and Mechanisms of Past Changes.” *Palaeogeography, Palaeoclimatology, Palaeoecology* 194(1–3):5–22.
- Glantz, M. 1996. “Currents of Change: El Niño’s Impact on Climate and Society.” Cambridge, U.K.: Cambridge University Press.
- Hall, G., and H. A. Patrinos. 2005. *Indigenous Peoples, Poverty and Human Development in Latin America, 1994–2004*. World Bank, Washington, DC.
- Halloy, S.R.P., Seimon A., Yager K. et al. 2005. “Multidimensional (Climate, Biodiversity, Socioeconomics, Agriculture) Context of Changes in Land Use in the Vilcanota Watershed, Peru.” In: *Land Use Changes and Mountain Biodiversity*, E.M. Spehn, M. Liberman Cruz, and C. Koerner, eds. CRC Press LLC, Ft. Lauderdale, pp. 323-337.
- IPCC (Intergovernmental Panel of Climate Change). 2007. “Climate Change 2007: The Physical Science

Basis.” Contribution of Working Group I to the Intergovernmental Panel of Climate Change, S. Solomon, D. Qin, M. Manning, Z. Chen, M. Marquis, K. B. Averyt, M. Tigner, and H. L. Miller, eds. Cambridge, U. K. and New York: Cambridge University Press.

IPCC (Intergovernmental Panel on Climate Change). 2001a. “Climate Change 2001: The Scientific Basis.” Contribution of the Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change, J. T. Houghton, Y. Ding, D. J. Griggs, M. Noguer, P. J. van der Linden, X. Dai, K. Maskell, and C. A. Johnson, eds. Cambridge, U.K.: Cambridge University Press.

IPCC (Intergovernmental Panel on Climate Change). 2001b. “Climate Change 2001: Impacts, Adaptation and Vulnerability.” A Contribution of the Working Group II to the Third Assessment Report of the Intergovernmental Panel on Climate Change, J. J. McCarthy, O. F. Canziani, N. A. Leary, D. J. Dokken, and K. S. White, eds. Cambridge, U.K.: Cambridge University Press.

IPCC (Intergovernmental Panel on Climate Change). 2007a. Climate Change 2007: The Physical Science Basis. Contribution of the Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change, ed. Salomon S., D. Qin, M. Manning, M. Marquis, K.B. Averyt, M. Tignor and H.L. Miller. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 996 pp.

IPCC (Intergovernmental Panel on Climate Change). 2007b. Climate Change 2007: Impacts, Adaptation and Vulnerability. Contribution of the Working Group II to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change, ed. Parry M.L., O.F. Canziani, J.P. Palutikof, P.J. van der Linden and C.E. Hanson. Cambridge University Press, Cambridge, UK, 976 pp.

IRI (International Research Institute for Climate and Society). 2001. *Coping with Climate—A Way Forward*. A multistakeholder review of regional climate outlook forums concluded at an international workshop. New York: IRI Publications.

Jordan, E. 1991. “Die Gletscher der bolivianischen Anden: eine photogrammetrische-kartographische Bestandsaufnahme der Gletscher Boliviens als Grundlage für klimatische Deutungen und Potential für die wirtschaftliche Nutzung”. Erdwissenschaftliche Forschung, Akademie der Wissenschaft und Literatur, Mainz, 23, Stuttgart.

Kaser, G. 2001. “Glacier–Climate Interaction at Low Latitudes.” *Journal of Glaciology* 47(157):195–204.

Orlove, B. S., J. C. H. Chiang, and M. A. Cane. 2000. “Forecasting Andean Rainfall and Crop Yield from the Influence of El Niño on Pleiades Visibility.” *Nature* 403:68–71.

Parmesan, C., and G. Yohe. 2003. A Globally Coherent Fingerprint of Climate Change Impacts Across Natural Systems.” *Nature* 421: 37–42.

Seimon, A. 2004. Unpublished Report.

Seimon, A. 2006. “Adaptation Strategies to the Environmental and Socioeconomic Impacts of El Niño for Rural Communities in Ecuador and Peru: Climatological and Meteorological Contexts.” Consultancy Paper to Climate Change Team, World Bank.

Sperling, F., and F. Szekely. 2005. “Disaster Risk Management in a Changing Climate.” Informal Discussion Paper prepared for the World Conference on Disaster Reduction on behalf of the Vulnerability and Adaptation Resource Group (VARG). Washington, D.C.


Stott, P.A., D.A. Stone and M.R. Allen. 2003. “Human Contribution to the European Heatwave in 2003” *Nature* 432: 610–614.


Valdivia, C., and R. Quiroz. 2006. “Adaptation Strategies to the Environmental and Socioeconomic Impacts of El Niño for Rural Communities in Ecuador and Peru.” Consultancy Report of Community Assessments in Peru conducted by Centro Internacional de la Papa in collaboration with ITDG and CIRNMA. Submission to the Climate Change Team, World Bank.


Valdivia, C., and J. Gilles. 2001. “Gender and Resource Management: Households and Groups,


- Strategies and Transitions.” *Journal of Agriculture and Human Values* 18(1):5–9.
- Valdivia, C. 2001. “Gender, livestock Assets, Resource Management, and Food Security: Lessons from the SR-CRSP.” *Journal of Agriculture and Human Values* 18(1):27–39.
- Valdivia, C., J. L. Gilles, C. Jetté, R. Quiroz, and R. Espejo. 2003. “Coping and Adapting to Climate Variability: The Role of Assets, Networks, Knowledge and Institutions.” In: *Insights and Tools for Adaptation: Learning from Climate Variability*, 189–199. Washington, D.C.: NOAA Office of Global Programs, Climate and Societal Interactions.
- Valdivia, C. 2004. “Andean Livelihoods and the Livestock Portfolio.” *Culture and Agriculture* 26(Spring 1&2):19–29.
- Vergara, V. 2005. “Adapting to Climate Change—Lessons Learned, Work in Progress, and Proposed Next Steps for the World Bank in Latin America.” Latin America and Caribbean Region Sustainable Development Working Paper 25. World Bank, Washington, DC.
- Vergara, W., Deeb. A.M., Valencia. A.M., Bradley R.S., Francou B., Zarzar A., Grünwaldt A., Haeussling S.M. 2007. “Economic Impacts of Rapad Glacier Retreta in the Andes” EOS, Transactions American Geophysical Union, Vol. 88, No. 25, 261-268.
- Vogel, C., and K. O’Brien. 2004. “Vulnerability to Global Environmental Change.” AVISO policy note, published by GECHS Project (www.gechs.org).
- Vuille, M. and R.S. Bradley. 2000. “Mean Annual Temperature Trends and Their Vertical Structure in the Tropical Andes”. *Geophysical Research Letters*, 27: 3885-3888.
- Vuille, M., R. S. Bradley, and F. Keimig. 2000. “Climate Variability in the Andes of Ecuador and Its Relation to Tropical Pacific and Atlantic Sea Surface Temperature Anomalies. *Journal of Climate* 13(14):2520–2535.
- Vuille, M., R. S. Bradley, M. Werner, and F. Keimig. 2003. “20th Century Climate Change in the Tropical Andes: Observations and Model Results.” *Climatic Change* 59(1–2):75–99.
- World Bank. 2005. “Natural Disaster Hotspots: A Global Risk Analysis.” Washington, DC: World Bank, 132 pp.
- World Bank. 2006. “Country Partnership Strategy for the Republic of Peru.” Washington, DC: World Bank.


Appendix A — Spatial Hazard Maps of Puno Communities











Appendix B — Warm (El Niño) and Cold (La Niña) Phases of ENSO

Year	DJF	JFM	FMA	MAM	AMJ	MJJ	JJA	JAS	ASO	SON	OND	NDJ	Year
1950	-1.8	-1.5	-1.4	-1.4	-1.4	-1.2	-0.9	-0.8	-0.8	-0.8	-0.9	-1	1950
1951	-1	-0.8	-0.6	-0.4	-0.2	0.1	0.4	0.5	0.6	0.7	0.7	0.6	1951
1952	0.3	0.1	0.1	0.1	0	-0.2	-0.3	-0.3	-0.1	-0.2	-0.2	-0.1	1952
1953	0.1	0.3	0.4	0.5	0.5	0.4	0.4	0.4	0.4	0.4	0.3	0.3	1953
1954	0.3	0.2	-0.1	-0.5	-0.7	-0.7	-0.8	-1	-1.1	-1.1	-1	-1	1954
1955	-1	-0.9	-0.9	-1	-1.1	-1	-1	-1	-1.5	-1.8	-2.1	-1.7	1955
1956	-1.2	-0.8	-0.7	-0.6	-0.6	-0.6	-0.7	-0.8	-0.9	-0.9	-0.9	-0.8	1956
1957	-0.5	-0.1	0.2	0.6	0.7	0.8	0.9	0.9	0.8	0.9	1.2	1.5	1957
1958	1.6	1.5	1.1	0.7	0.5	0.5	0.4	0.1	0	0	0.1	0.3	1958
1959	0.4	0.4	0.3	0.2	0	-0.3	-0.4	-0.5	-0.4	-0.4	-0.3	-0.3	1959
1960	-0.3	-0.3	-0.3	-0.2	-0.1	-0.1	0	0	-0.1	-0.2	-0.3	-0.2	1960
1961	-0.2	-0.2	-0.2	-0.1	0.1	0.1	0	-0.3	-0.6	-0.6	-0.5	-0.5	1961
1962	-0.5	-0.5	-0.5	-0.5	-0.4	-0.3	-0.2	-0.3	-0.4	-0.6	-0.7	-0.7	1962
1963	-0.6	-0.3	0	0.1	0.1	0.3	0.6	0.8	0.8	0.9	1	1	1963
1964	0.8	0.4	-0.1	-0.5	-0.7	-0.7	-0.8	-0.9	-1	-1.1	-1.1	-1	1964
1965	-0.8	-0.5	-0.3	0	0.2	0.6	1	1.2	1.4	1.5	1.6	1.5	1965
1966	1.2	1.1	0.8	0.5	0.2	0.1	0.1	0	-0.2	-0.3	-0.3	-0.4	1966
1967	-0.4	-0.5	-0.6	-0.5	-0.3	0	0	-0.2	-0.4	-0.5	-0.5	-0.6	1967
1968	-0.7	-0.9	-0.8	-0.8	-0.4	0	0.3	0.3	0.2	0.4	0.6	0.9	1968
1969	1	1	0.9	0.7	0.6	0.4	0.4	0.4	0.6	0.7	0.7	0.6	1969
1970	0.5	0.3	0.2	0.1	-0.1	-0.4	-0.6	-0.8	-0.8	-0.8	-0.9	-1.2	1970
1971	-1.4	-1.4	-1.2	-1	-0.8	-0.8	-0.8	-0.8	-0.9	-0.9	-1	-0.9	1971
1972	-0.7	-0.3	0	0.3	0.5	0.8	1.1	1.3	1.5	1.8	2	2.1	1972
1973	1.8	1.2	0.5	-0.1	-0.5	-0.8	-1.1	-1.3	-1.4	-1.7	-1.9	-2	1973
1974	-1.8	-1.6	-1.2	-1.1	-0.9	-0.7	-0.5	-0.4	-0.5	-0.7	-0.8	-0.7	1974
1975	-0.6	-0.6	-0.7	-0.8	-1	-1.1	-1.3	-1.4	-1.6	-1.6	-1.7	-1.8	1975
1976	-1.6	-1.2	-0.9	-0.7	-0.5	-0.2	0.1	0.3	0.5	0.7	0.8	0.8	1976
1977	0.6	0.5	0.2	0.1	0.2	0.3	0.3	0.4	0.5	0.7	0.8	0.8	1977
1978	0.7	0.4	0	-0.3	-0.4	-0.3	-0.4	-0.5	-0.5	-0.4	-0.2	-0.1	1978
1979	-0.1	0	0.1	0.2	0.1	0	0	0.2	0.3	0.4	0.5	0.5	1979
1980	0.5	0.3	0.2	0.2	0.3	0.3	0.2	0	-0.1	0	0	-0.1	1980
1981	-0.3	-0.4	-0.4	-0.3	-0.3	-0.3	-0.4	-0.3	-0.2	-0.1	-0.1	-0.1	1981
1982	0	0.1	0.2	0.4	0.6	0.7	0.8	1	1.5	1.9	2.2	2.3	1982
1983	2.3	2	1.6	1.2	1	0.6	0.2	-0.2	-0.5	-0.8	-0.9	-0.8	1983
1984	-0.5	-0.3	-0.2	-0.4	-0.5	-0.5	-0.3	-0.2	-0.3	-0.6	-1	-1.1	1984
1985	-1	-0.8	-0.8	-0.8	-0.7	-0.5	-0.4	-0.4	-0.4	-0.3	-0.2	-0.3	1985
1986	-0.4	-0.4	-0.3	-0.2	-0.1	0	0.2	0.5	0.7	0.9	1.1	1.2	1986
1987	1.3	1.2	1.1	1	1	1.2	1.5	1.6	1.6	1.5	1.3	1.1	1987
1988	0.8	0.5	0.1	-0.3	-0.8	-1.2	-1.2	-1.1	-1.3	-1.6	-1.9	-1.9	1988
1989	-1.7	-1.5	-1.1	-0.9	-0.6	-0.4	-0.3	-0.3	-0.3	-0.3	-0.2	-0.1	1989
1990	0.1	0.2	0.3	0.3	0.3	0.3	0.3	0.4	0.3	0.3	0.3	0.4	1990
1991	0.5	0.4	0.4	0.4	0.6	0.8	0.9	0.9	0.8	1	1.4	1.7	1991
1992	1.8	1.7	1.6	1.4	1.1	0.8	0.4	0.2	-0.1	-0.1	0	0.1	1992
1993	0.3	0.4	0.6	0.8	0.8	0.7	0.5	0.4	0.4	0.3	0.2	0.2	1993
1994	0.2	0.3	0.4	0.5	0.6	0.6	0.6	0.6	0.7	0.9	1.2	1.3	1994
1995	1.2	0.9	0.7	0.4	0.2	0.1	0	-0.3	-0.5	-0.6	-0.7	-0.8	1995
1996	-0.8	-0.7	-0.5	-0.3	-0.2	-0.2	-0.1	-0.2	-0.2	-0.2	-0.3	-0.4	1996
1997	-0.4	-0.3	0	0.4	0.9	1.4	1.7	2	2.3	2.4	2.5	2.5	1997
1998	2.4	2	1.4	1.1	0.4	-0.1	-0.8	-1	-1.1	-1.1	-1.3	-1.5	1998
1999	-1.6	-1.2	-0.9	-0.7	-0.8	-0.8	-0.9	-0.9	-1	-1.2	-1.4	-1.6	1999
2000	-1.6	-1.5	-1.1	-0.9	-0.7	-0.6	-0.4	-0.3	-0.4	-0.5	-0.7	-0.7	2000
2001	-0.7	-0.5	-0.4	-0.2	-0.1	0.1	0.2	0.1	0	-0.1	-0.2	-0.2	2001
2002	-0.1	0.1	0.3	0.4	0.7	0.8	0.9	0.9	1.1	1.3	1.5	1.3	2002
2003	1.1	0.8	0.6	0.1	-0.1	0	0.3	0.4	0.5	0.5	0.6	0.5	2003
2004	0.4	0.2	0.2	0.2	0.3	0.4	0.7	0.8	0.9	0.9	0.9	0.8	2004
2005	0.6	0.5	0.3	0.4	0.5	0.3	0.2	0	0	-0.2	-0.4	-0.7	2005
2006	-0.8	-0.7	-0.4	-0.2	0	0.1	0.3	0.4	0.7	0.9	1.1	1.1	2006
2007	0.8	0.3	0.1	-0.1	0	-0.1	-0.2	-0.6					2007

Definition: Conditions for a warm (cold) phase of ENSO are met if SST anomalies above 5°C (below -5°C) are sustained for a minimum of three consecutive months.


Source: NOAA http://www.cpc.ncep.noaa.gov/products/analysis_monitoring/ensostuff/ensoyears.shtml.

Appendix C — Precipitation Totals in Relation to Seasonal SST Anomalies in Esperanza


Seasonal precipitation totals in relation to seasonal SST anomalies in the NINO 3.4 region in Esperanza, coastal northwestern Peru

The graphs show examples of different lead times for SST anomalies in the NINO 3.4 region in relation to the later observed precipitation total measured for Esperanza from January to March (JFM).


SST Anomalies from May to July (MJJ) of the Preceding Year




SST Anomalies from June to August (JJA) of the Preceding Year




SST Anomalies from August to September (ASO) of the Preceding Year




SST Anomalies from October to December (SON) of the Preceding Year



SST Anomalies from November to January (NDJ)



SST Anomalies from December to February (DJF)



Appendix D — Community-Specific Recommendations

Parachique

When the groups were asked about future measures that the community and its families could implement to strengthen their ability to adapt and respond to climate events, the responses from the men's and women's groups were different. The men mentioned policy proposals that would let them manage the resources that they generate for developing the fishing community (an expression of the situation of conflict with the Municipality of Sechura and with central government organizations, by whom they feel ignored), support for improving artisanal fishing activities, making the bay healthier, coastal defense, and organizing and training the population. The women proposed measures at the family and daily level (they expressed a great deal of concern during the workshop about the safety conditions in their homes and in the bay).

Measures that Do Not Require Outside Assistance

Measures proposed by the men's group:

- Continue with the climate information provided by the Parachique Harbormaster's Office, because it is reliable.
- Comply with the Parachique Harbormaster Office's permission to leave port for any boat heading out to sea.
- Involve the entire family, including women and young children, in discussions about the future of Parachique.

Measures proposed by the women's group:

- Believe in religion, which helps people to cope with disasters.
- Secure homes well to keep water out.

Measures that Require Outside Assistance

Measures proposed by the men's group:

Work for legislation and standards that favor the town's desires

- Get the community recognized as a Minor Population Center (Centro Poblado Menor) with represented municipality functions, so that its tax contributions to the Municipality of Sechura are reinvested in Parachique.
- Surveillance and compliance with the laws that protect artisanal fishing.
- Get the Ministry of Fisheries to return the amount of taxes collected from the community, to be used for reinforcing the seawall.

Support the improvement of artisanal fishing activities

- Create a fund to finance the improvement and updating of fishing boats and methods, so that fishermen can also catch the new species that appear with El Niño, and finance navigation and radio communication equipment for the boats.
- Promote diversification between artisanal fishing and other activities.
- Support the installation of a refrigeration chamber to preserve artisanal fishing products.

- Get practical technical training related to fishing activities.

Improve health conditions in the bay

- Improve the area's health conditions in order to protect the people's health and improve fishing production (they don't have health certification for the fishing products).

Protection, safety, information

- Improve and expand coastal defense.
- Install a communications system for direct warnings between the Callao Harbormaster's Office and the Parachique Harbormaster's Office in case of ocean phenomena.
- Establish weather stations in Parachique, especially for reading ocean temperature and measuring winds.
- Support the installation of a hyperbaric chamber for rapid first aid to shellfish divers.
- Establish a climate change office in Parachique (they say there are young university students who could staff it).

Organization and training

- Support better organization of the group that represents the entire town.
- Conduct massive training workshops to promote the population's awareness and consideration regarding its attitude toward weather phenomena.

Communications equipment

- Install a public telephone for receiving calls and using the Internet.
- Install a parabolic antenna for retransmitting TV broadcasts.

Measures proposed by the women's group:

- Have family vegetable gardens.

- Build homes in the higher section for those that do not have them.
- Develop an adult education school.

Chato Grande

The participants of each group were asked about future measures that the community and its families could implement to strengthen their ability to adapt and respond to climate events. They identified the following:

Measures the Community Could Take without Outside Assistance

Water Management

- Generate agreements with the Irrigation Users Committee to improve water management;
- Develop crops that require little water, such as black-eyed peas and corn; and
- Greater concern for and monitoring of irrigation water turns.

Social Participation

- Involve young people and teachers in communal work and efforts.

Preventive Evacuation

- Move to Nuevo Chato Grande sufficiently in advance. This evidently involves a "learned lesson," since they have no actual plans to move because they consider Nuevo Chato Grande too distant from their income sources.

Measures that Require Outside Assistance

Flood-risk mitigation

- Improve river-bank defenses by raising the levees.
- Dredge the river to remove the silt.
- Change the river's course and provide it with an outlet into the sea (14 km) with the involvement

of the central government (Ministry of Agriculture) and the regional government. This is a deeply felt need; the participants said it could be done, because they have heard proposals in this regard; but machinery and a major investment are required.

Water and environmental management

- Improve management of irrigation and human-consumption water management.
- Train residents and farmers on water management and use vis-à-vis major drought issues.
- Protect the forest by planting more trees and controlling logging.
- Promote a fair distribution of irrigation water.
- Reinforce environmental education at the school level.

Diversifying of activities

- Develop training centers and workshops for technical works such as plumbing, masonry, etc.
- Promote forming of microbusinesses for income generation.
- Seek, in an organized manner, income-generating alternatives for drought periods (pottery, weaving, beekeeping).

Improvement of agriculture, raising livestock, and promotion of processing activities

- Promote and support production of profitable crops of short vegetative periods that require little water (such as peanuts, soybeans, black-eyed peas, pigeon peas, grapes, watermelon, and squash).
- Improve pig raising.
- Provide technical advisory, training, and credit for the development of agroforestry systems to protect crops.
- Promote the development of activities that add value to local products and further employment; for instance, carob-processing activities.

Infrastructure improvement

- Improve the road between Nuevo Chato Grande and Chato Grande Antiguo (Old Chato Grande).

Preparation and prevention for El Niño effects and climatic variability

- Create a family fund to cope with El Niño events.
- Train and organize the population for preparedness and better coping with contingencies.
- Provide access to clear and easily understandable climate information to aid the farmers' decision making, including practical advice for their planting decisions.
- Access to timely information to issue flood warnings on time.
- Build the local promoters' capacities. This proposal emerged from the group's discussion as a reflection on the need to have a preventive attitude and to encourage the population to adopt changes.

Locuto

The participants were asked about future measures that the community and its families could implement to strengthen their ability to adapt and respond to climate events. They identified the following:

Measures the Community Could Take without External Assistance

- Sensitize the population (especially children) about climate changes and El Niño.
- Strengthen the community's ability to form and maintain organization.
- Promote recognition of women's value in the community.
- Take advantage of local resources in production and agribusiness.
- Promote diversification of forest-related activities.
- Promote growing of tamarind and other species that require little water.

- Each family could plant two tamarind plants on their land now, which would start producing in three to four years.
- In the future, select a variety of seeds for reforestation: carob, overal, sapote.
- Promote changes in eating habits that increase consumption of local products (carob coffee, carob powder, and milk consumption).

Measures that Require Outside Assistance

- Improve education quality.
- Investigate possible low-risk production on land in the dry forest area, using well water or pumped river water.
- Investigate the use of subsoil water and tubular wells for human consumption on the left riverbank.
- Organize and train the population for preparations and prevention.
- Prepare a rational forest management plan.
- Develop a genetic improvement plan for sheep to be used for meat.
- Develop organic beekeeping methods using the virgin forests.
- They are currently working with Plan Piura (formerly Plan Internacional) to develop organic fertilizers and better stoves, and the educational centers are conducting projects to process carob products. Cofinancing and management are needed from the regional and central governments and from nongovernmental organizations.

The community of Locuto is part of an area called the left bank of the Piura River in Tambogrande. In this area and in others within the district, they recently began a participatory planning process to formulate a development plan, directed by the Municipality of Tambogrande and with technical support from the Ceproda Minga and Cooperación nongovernmental organizations. The measures discussed, identified in a workshop held on July 13, 2005 became the framework

of the proposed development plan for the area and are listed below:

Measures that Could Be Implemented by the Community without External Assistance (part of the development plan for the left bank of the Piura River, in Tambogrande)

- Storage of carob for feeding livestock.
- Coordination with SENASA, local government, and nongovernmental organizations to run livestock disease prevention campaigns.
- Improvement of livestock pens.
- Organization to implement production initiatives.
- Improved strategies for aiding the most vulnerable families.
- Diversified production and income-generating activities.
- Tree thinning and pruning as part of a dry-forest management strategy.
- Communitywide control measures to reduce logging and charcoal production.
- Make the decision to relocate the homes of families affected by the 1997–98 El Niño.
- Plan to plant the occasional and floodplain fields as a community, promoting diversification with resistant varieties and reducing dependency on the use of agrochemicals.
- Agree not to increase the areas intended for use as occasional fields, promoting diversification of production activities.
- Encourage the consumption of yupisin to combat anemia and malnutrition.
- Improve corrals to protect livestock.
- Encourage and promote the population's participation through these organizations.
- Work with public and private institutions.
- Hold meetings to identify and teach the lessons learned during the 1982–83 and 1997–98 El Niño events.

Measures that Require External Assistance (part of the development plan for the left bank of the Piura River, in Tambogrande)

Planning and regulation

- Formulate a territorial organization plan, with participation and with a focus on prevention, to divide the forest into zones based on the local population's production, extraction, and reproduction activities; use it as the planning unit for the entire dry-forest ecosystem on the left bank of the Piura River, with jurisdiction in the Districts of Tambogrande and Chulucanas.
- With participation, develop a dry forest management plan covering the four farming communities on the left bank of the Piura River, Tambogrande District.
- Develop a forest fire prevention and control plan.
- Design an organizational plan of areas to be used for housing and services.

Technical studies and projects

- Conduct a technical study on risk evaluation and assignment by territory.
- Conduct the technical study and construct a potable water system for the left bank of the Piura River in the District of Tambogrande.
- Conduct a technical study on soil conditions in the current occasional field areas to determine the level of degradation, potential use, and recovery efforts needed, as well as low-risk production possibilities using subsoil water.
- Complete the technical study for building a bridge over the Piura River.
- Drill community-chain pump wells to supply water for family consumption and to improve productive activities such as raising small livestock.
- Build firebreak ditches in areas where forest fires are common.
- Construct a drainage system in inhabited areas to accommodate the rainy seasons.

- Execute projects and take measures to control watercourses and gullies with appropriate techniques using local resources.

Improvement of rural production systems

- Control carob pests and diseases using methods based on ecological principles and management that reinforces the natural vegetation's diversity.
- Conduct research on the diversity of pests that affect carob flowering and fruit production, given the importance of the carob pods in forest-production activities.
- Promote the construction of family carob silos.
- Promote rural agribusiness and diversification of production activities based on proper management of the dry forest.
- Establish a permanent community animal health program.
- Conduct genetic improvement in sheep to use for meat.
- Introduce tamarind, a variety that is well adapted to dry forest conditions and for which there is a market demand.
- Create a seed bank with varieties adapted to El Niño conditions for planting in the occasional and floodplain fields.
- Conduct training, credit, technical-assistance, and market-access programs to improve beekeeping and rural agribusiness and to promote agroecological production.
- Certify products as organic.
- Promote ecological tourism.

Nutrition, health, and social aspects

- In all municipalities, institute the procurement of soluble carob powder to supplement the Glass of Milk Program's allotments.
- Promote the construction of better stoves to reduce wood consumption and improve women's health conditions.

- Implement nutritional support and food-for-work programs for highly vulnerable families in drought situations.
- Institute a program for disease prevention and for treating the symptoms and aftereffects of malaria.

Organizational strengthening, capacity building, and education

- Strengthening the Locuto community's social organizations, developing prevention, and response capabilities.
- Build the organizational capacities of Locuto's Apóstol Juan Bautista Peasant Community and the settlements it comprises, promoting recognition of women's value in society.
- Support the promotion of leadership and reinforcement in productive and commercial organizations.
- Implement permanent environmental education programs, with a preventive focus, that promote sustainable management of the dry forest.

Response organization and preparation

- Implementation of a community early warning system, incorporating local knowledge.
- Training and equipment for civil defense committees.
- Adequate and timely access to weather forecasts.

Cabuyal

The participants were asked about future measures that the community and its families could implement to strengthen their ability to adapt and respond to climate events. They identified the following:

Measures the Community Could Take without Outside Assistance

Improvement of farming-production systems

- Recover and spread cultivation of native food crops (such as the pajuro and vitalicio bean, etc.) that were substituted by introduced species.

- Change planting dates according to climate conditions to reduce risks.

Improvement of livestock production systems

- Processing of harvest residues to feed livestock (using bagasse and stubble in livestock feed).
- Develop a sanitary calendar to control livestock.

Resource management and conservation

- Perform soil conservation practices such as the construction of hillside terraces, fallowing, and crop rotation.
- Reforestation with native species in gullies, such as planting sangoyuyo to protect water sources.

Education, information, organization, and social practices

- Promote sharing information with farmers from other places.
- Recover and strengthen traditional social, economic, and cultural networks and practices that contribute to the risk-reduction strategies in the community and the families (for instance, save the seed in the Andean plateau, as a protection strategy against extreme weather events).

Production Infrastructure

- Improve irrigation systems by strip cropping (the field is divided into small 5-meter strips crosswise to the slope of the field and ditches that run parallel to each strip are dug to irrigate the crops).

Measures that Require Outside Assistance

Improvement of farm-production systems

- Promote growing crops that are profitable crops and less sensitive to climate variability (resistant to water excess and shortage) such as sugarcane, pigeon peas, etc.

- Recover the use of true potato seed to produce potato for consumption and seed potato, as a strategy for quick food production under crisis situations; for which native potato seeds could be bought in Huancabamba and in the Andean plateau.
- Facilitate market access to improve the price of agricultural and livestock products.
- Disseminate agroecological farming practices, such as the use of organic fertilizers and pesticides.
- Plan the agricultural calendar using local knowledge and scientific forecasts.
- Recover traditional knowledge on climate; study its indicators and accuracy level.

Improvement of livestock-production systems

- Improve techniques for livestock-related activities, through changes in extensive and intensive (stalls) raising systems, and planning livestock development based on the load capacity of the land.
- Improve livestock (meat and milk).
- Appropriately manage livestock and pastures.
- Request technical assistance to governmental agencies (Ministry of Agriculture).
- Perform sanitary control of livestock (Tupe in cows).

Education, information, organization, and social practices

- Promote further sensitizing of the population in regard to climate change.
- Train the population on disaster prevention and response.
- Strengthen social organizations to recover and build their capacity for collective action (solidarity, reciprocity, and community work practices) and management.
- Establish a link between the school and the major issues and aspirations of the community.

Social infrastructure

- Plan the housing sites where the houses will be located to reduce the risk of houses collapsing due to erosion and activation of watercourses during heavy rains.
- Perform technical studies to ensure a safer site for the potable water system.

Production infrastructure

- Improve and protect the irrigation system, diminishing the extent of infiltration and its vulnerability to landslides.
- Make studies on the water supply of the Ñoma River available to the population, to use them as a basis for crop planning.
- Upgrade and equip the panela sugarcane mills.

Appendix E — Project Proposal for Follow-Up Activities in the Altiplano

Background

The Bolivian and Peruvian altiplano is among the regions in Latin America with the highest incidence of extreme and persistent rural poverty. In a highly variable climate, livelihoods based on farming practices are consistently challenged. Frequent extreme events, including droughts, floods, hail, and cold spells make it difficult for households to accumulate assets and often reinforce conditions of poverty. From 1970–1999, estimates for Bolivia suggest that total accumulated economic losses associated with natural disasters add up to 1 percent of annual GDP. In Peru, extreme poverty is especially widespread in the rural areas of the highlands and among the indigenous population. This is particularly striking, as according to INDECI, Puno in the Peruvian altiplano ranked highest among all departments in terms of emergencies reported from 2000 to 2004 linked to climatic hazards. Research by Valdivia, Quiroz and others has shown that prolonged and repeated climatic shocks have triggered migration in Bolivia and Peru, transposing poverty to urban areas.

This vulnerability is being exacerbated by environmental and socioeconomic changes. Mountain ecosystems are particularly exposed to climate change (IPCC 2001), as the effects of global warming are more pronounced at high altitudes. Significant warming trends are already visible in the Andes, raising concerns about the effects on ecosystems and water resources, and associated downstream effects for agricultural activities in the altiplano, which are already under pressure from environmental degradation.

These changes in physical conditions are becoming visible while economic pressures have reduced the diversity of agroecosystems (Valdivia, personal communication)—making them more vulnerable to environmental change. In particular, the largely indigenous communities (Quechua and Aymara) in the altiplano are faced with the challenging task of adapting their traditional livelihoods to these tremendous socioeconomic and environment pressures.

Development Objective

The objective of the project is to increase the resilience of rural livelihoods to climatic shocks by responding directly to capacity-building needs expressed in drawing on participatory assessments conducted by the World Bank, University of Missouri, Center for International Potato Research (CIP), local nongovernmental organizations, and others in the altiplano of Bolivia and Peru. In accordance with these identified needs, indigenous communities will receive training in natural resource management and agricultural practices best suited for the climate characteristics of the region. This includes education about environmentally suitable crops, planting techniques and pest management, markets for sustainable crops, and institutional support mechanisms. The content of the training will be informed by information from seasonal climate forecasts and spatial hazard-risk mapping for various climatic hazards, incorporating also historical experiences and knowledge of the communities. The proposed project will develop training initiatives and expand these to communities in the highlands of

Bolivia, as the exposure and vulnerabilities to climatic shocks in the Bolivian altiplano largely mirror the challenges faced in Peru and hence allow for a broader regional focus.

Expected Contribution to Development Outcomes/Results

The seventh Millennium Development Goals acknowledge the importance of environmental sustainability to the overarching objective of poverty alleviation. In a 10-agency report, “Poverty and Climate Change,” led by the World Bank, the importance of addressing current and future climate risks for sustaining progress toward and beyond the MDGs. The World Conference on Disaster Reduction and the United Nations Framework Convention on Climate Change called upon the international community to address the links between disaster risk management, adaptation to climate change, and development.

This link between development and environment is particularly evident in the altiplano, where livelihoods are threatened by natural hazards and environmental degradation. The forthcoming Country Environmental Assessment for Peru highlights the high vulnerability of the altiplano to disaster risks and emphasizes the general need for improving the risk management framework. Climate-change projections further reinforce this picture by exacerbating the pressure on natural resources.

By developing practical risk management options for rural livelihoods in the altiplano, the project addresses community-level capacity-building needs, while generally contributing to an improved practical understanding of how the resilience of development processes to exogenous shocks and global change can be improved.

Expected Outputs and Results

The key output, which directly aims at improving development prospects, is the provision of practical training to communities for increasing the resilience to climatic shocks. The goal is to train a minimum of six communities in the Peruvian altiplano and in Bolivia. The project will consolidate local knowledge networks and partnerships initiated during a previous World Bank trust-funded project in Peru and the research activities of our collaborating partners in Bolivia.

The training will provide linkages between producer and expert knowledge systems; it is therefore conducted as a highly interactive process between researchers and communities. For wider regional use and replication of the approach, the following documents will be produced: i) Methodology on participatory hazard-risk mapping, ii) methodology on converting climate forecasts into information formats relevant to rural Andean communities, iii) report on risk-mitigating options in natural resource management and agricultural practices, iv) crossregional workshop (intended for optional third year of project) discussing natural resource management in marginal regions in Latin America, Africa, and Asia (focusing on countries with related environmental challenges; envisioned target countries include Ethiopia, Eritrea, Bhutan, and Mongolia), and v) report on lessons learnt for building comprehensive risk-management strategies for rural livelihoods exposed to natural hazards and environmental change (with an emphasis on Latin America, but with a crosscomparison with projects in other regions, drawing on workshop input).

Key Performance Indicators

Quantifiable indicators include i) the number of communities trained (with the objective of improving linkages between producer and expert-knowledge systems, which promote the application of risk-

mitigating strategies by rural households), ii) practical guidance (reports and briefing notes) to the World Bank and other development agencies supporting the integration of risk mitigation into rural development

processes, and iii) (long-term) improved resilience of targeted communities to climatic shocks, exemplified by the ability to accumulate and develop assets, and the reduced number of household-level disasters.


Appendix F — Managing Climate Risks in Coastal Ecuador — A Brief Comparison of Community Perspectives

Coastal Ecuador's exposure to natural hazard is similar to the Northwestern part of Peru. The El Niño Southern Oscillation (ENSO) represents the dominant influence of inter-annual climate variability. During El Niño events, the peak monthly rainfall amount increases several fold in comparison to normal years and the rainy season tends to be longer (see Figure). In addition, El Niño is associated with extreme rainfall events that trigger destructive floods. The ENSO influence is superimposed on a marked intra-seasonal variability in coastal Ecuador. The amplitude in variability increases from North to South with a distinct rainy and dry season.

Community assessments led by Jorge Marcos and Pilar Cornejo of Escuela Superior Politécnica del Litoral (ESPOL), covering eight communities in the Southern Manabi Province, Guayas Province and El Oro Province, yielded results which are largely comparable to the ones uncovered in Peru.

As in Piura, Peru, the communities have a clear understanding of El Niño, but not of La Niña. Communities associate El Niño with excessive rains and destructive floods. Only strong El Niño events are recalled, as weak or moderate events are not being distinguished from a strong rainy season. Hence, the

Figure. Monthly Precipitation Averages and Totals Observed During El Niño years, Guayaquil, Ecuador



Note: The figure shows how monthly precipitation totals generally increase significantly and the rainy season lengthens during El Niño years (i.e., 1982/83, 1986/97, 1997/98) in comparison to the long-term monthly averages observed over 30+ years in Guayaquil (sea level, Latitude: 02°09', Longitude: 79°53'W). While El Niño events magnify the rainy season, each event has its own distinct characteristics depending in part on the timing of the onset and cessation relative to seasonal patterns.

period between individual El Niño events in the eyes of the communities is longer than the return period of El Niño events, when using scientific classifications.

There is no clear understanding of La Niña as a concept. In fact the term can be misleading. La Niña is sometimes understood by community members as a more extreme El Niño. As a female version of El Niño (“the boy child”), it is associated with more “crying” and hence more rainfall. This interpretation runs contrary to the scientific definition, where La Niña represents the cold phase of ENSO and is linked to drier than normal conditions in coastal Ecuador. In addition, there was a wide-spread perception among community members that the diurnal temperature cycle has become more extreme and the seasons have become less reliable.

Communities are particularly vulnerable to climate related impacts on human health, infrastructure and agriculture. El Niño events are associated with an increase in water and vector borne diseases, such as malaria, dengue and cholera. Given the pervasive absence of adequate health facilities, returning migrant works represent an additional health risk. Floods triggered by excessive rainfall events have led to destruction of houses, roads and bridges. Communities highlighted the difficulty of accessing the markets during El Niño events and years with strong rainfall. This constrains the coping options of households, as they cannot sell livestock or acquire new seeds. The loss of infrastructure has also interrupted of schooling time of their children. Agriculture and livestock during excessively wet periods is affected through destruction of planting sites, increase in plant pests and animal diseases. Medium-term effects of El Niño for agriculture can be positive as the environment is revitalized and planting areas can be expanded.

Given the heavy dependency of livelihoods on agricultural activities, failed or erratic rainy seasons lead to the depletion of assets or force members of households to migrate to urban centers in search of

labor. There was a wide-spread perception among communities that the diurnal temperature cycle has become more extreme over recent years and the rainy season less reliable. Where access to irrigation is not available, traditional water collection structures can help reduce vulnerabilities against drought conditions.

In contrast to the assessments in Piura, traditional water storage structures, known as Albarradas or Jagüeyes are used by communities to minimize the effects of floods and droughts. Albarradas are u-shape detention ponds, which are build on semi-permeable surfaces. During rainfall events, run-off collects in these structures and recharges the aquifer, which can then be tapped during dry years through the construction of dry wells. Since the water percolates through the soil and collects below ground, evaporation rates are reduced. As a result the excessive rainfall during strong El Niño years is turned into an asset, helping communities to manage the dry periods.

There is archeological evidence that Albarradas have been constructed by the local populations as an adaptation strategy for over 3000 years. While Albarradas were in use in several of the communities visited in Ecuador, no such structures were discovered in the community assessments in Peru. The reasons for this are unclear, but may be linked to the different country histories and land-use policies. As the knowledge of the maintenance and with it the use of these ancient human-made structures is fading, it may be worthwhile to consider whether this trend should be reversed and examined in the broader regional applicability as a community level alternative to technological interventions for managing climate variability.

The scope of recommendations, which emerged from the community meetings and discussions with government stakeholders, echo the issues of the Peru assessments. These include (i) the improvement of climate information formats through a stronger engagement of communities and recognition of local

knowledge systems, (ii) strengthening awareness of health hazards in relation to climatic events and improving access to preventive measures, (iii) support the maintenance of water harvesting and storage structures, (iv) research and training on natural resource management, (v) sustainable use of the tropical dry forest, and (vi) improved access to credit, where payment schemes take into account the timing of agricultural practices.

A range of regional and national institutions, government ministries and universities are engaged in research and policy initiatives aimed at strengthening climate risk management efforts in Ecuador, including the Centro Internacional de Investigaciones de

Fenomeno del El Niño (CIIFEN), Escuela Superior Politecnica del Litoral (ESPOL), Instituto Nacional de Meteorología e Hidrología (INAMHI), Instituto Nacional de Pesca (INP), Ministerio de Agricultura and Ministerio del Ambiente. The recommendations of the community assessments are suited for a range of community driven development (CDD) or micro-projects. These could be used as a vehicle to further strengthen the linkage between scientific advances, institutional awareness, and local needs.

This brief summary is based on submissions by, and personal communication with, J. Marcos and P. Cornejo, ESPOL.



Environment Department
THE WORLD BANK

1818 H Street, NW
Washington, D.C. 20433
Telephone: 202-473-3641
Facsimile: 202-477-0565

